



DNA ENCODING SNORF62 AND SNORF72 RECEPTORS

5

BACKGROUND OF THE INVENTION

10 This application is a continuation-in-part of U.S. Serial  
15 No. 09/558,099, filed April 25, 2000 which is a  
20 continuation-in-part of U.S. Serial No. 09/466,435, filed  
December 17, 1999, the contents of which are hereby  
incorporated by reference into the subject application.

25 Throughout this application various publications are  
referred to by partial citations within parenthesis. Full  
citations for these publications may be found at the end of  
the specification immediately preceding the claims. The  
disclosures of these publications, in their entirety, are  
hereby incorporated by reference into this application in  
order to more fully describe the state of the art to which  
the invention pertains.

30 Neuroregulators comprise a diverse group of natural products  
that subserve or modulate communication in the nervous  
system. They include, but are not limited to,  
35 neuropeptides, amino acids, biogenic amines, lipids and  
lipid metabolites, and other metabolic byproducts. Many of  
these neuroregulator substances interact with specific cell  
surface receptors which transduce signals from the outside  
to the inside of the cell. G-protein coupled receptors  
(GPCRs) represent a major class of cell surface receptors  
with which many neurotransmitters interact to mediate their  
effects. GPCRs are characterized by seven membrane-spanning  
domains and are coupled to their effectors via G-proteins  
linking receptor activation with intracellular biochemical  
sequelae such as stimulation of adenylyl cyclase. This  
application describes the identification of two GPCRs,

SNORF62 and SNORF72, as receptors for neuromedin U (NMU) neuropeptides.

Neuropeptides are synthesized and released from neurons to mediate their effects on cells within the nervous system or on peripheral targets. NMU-25 and NMU-8 are bioactive peptides originally isolated from porcine spinal cord (Minamino, N. et al. 1985a and 1985b). NMU-8 corresponds to the C-terminus of porcine NMU-25 preceded by Arg-Arg residues and may therefore be generated by enzymatic cleavage. NMU homologues have been identified in many species including human (25 amino acids) and rat (23 amino acids).

The amino acid sequence for human NMU-25 is as follows:

F R V D E E F Q S P F A S Q S R G Y F L F R P R N-NH<sub>2</sub>  
(SEQ ID NO: 5).

The amino acid sequence for porcine NMU-25 is as follows:

F K V D E E F Q G P I V S Q N R R Y F L F R P R N-NH<sub>2</sub>  
(SEQ ID NO: 6).

The amino acid sequence for rat NMU-23 is as follows:

Y K V N E - Y Q G P - V A P S G G F F L F R P R N-NH<sub>2</sub>  
(SEQ ID NO: 7) (- indicates gaps in rat NMU-23 sequence to demonstrate optimum alignment).

The amino acid sequence for NMU-8 is Y F L F R P R N-NH<sub>2</sub> (SEQ ID NO: 8). All of the preceding sequences were taken from Nandha and Bloom 1993 and Austin, et al. 1995.

Interestingly, the 8 carboxy-terminal residues of human NMU-25 are identical to those in porcine, rabbit and guinea pig NMU and differ only by one residue from the C-terminus of frog, rat, dog and chicken NMU (Austin et al. 1995). C-terminal NMU peptides (8 - 9 amino acids) have also been

identified in guinea pig, chicken and dog tissue extracts (Minamino et al. 1985a and 1985b, Domin et al. 1989, O'Harte et al. 1991). Indeed, the region of rat NMU-23 critical for smooth muscle contractile activity was found to reside between residues 17 - 22 (C-terminal region) (Hashimoto et al. 1991, Sakura et al. 1991). However, other groups have demonstrated the necessity of the amidated C-terminal asparagine (Asn23) for activity as well (Nandha and Bloom 1993). Full length NMU is approximately 3-fold more potent than NMU-8 in smooth muscle contraction assays suggesting that the N-terminal region of the peptide also contributes to the activity (Nandha and Bloom, 1993). Several residues in the middle region of the peptide are conserved between species including Glu5, Gln8 and Pro10 supporting the functional importance of this region of the peptide (Nandha and Bloom, 1993). The C-terminus of NMU shares some homology with rat pancreatic polypeptide (PP): Leu-X-Arg-Pro-Arg-X-amide and contains a terminal asparaginamide also present in vasoactive intestinal polypeptide (VIP) (Nandha and Bloom 1993). However, the structure of NMU is unrelated to the other neuromedin peptides isolated by Minamino et al. (1985a,b).

A profound effect of NMU has been observed in rats on the *in vivo* release of stress-related modulators from the anterior pituitary and adrenal glands (reviewed in Malendowicz and Markowska 1994). Following a single subcutaneous injection of NMU-8 (6 µg/ 100g body weight), adrenocorticotrophic hormone (ACTH) blood concentrations are elevated transiently (3 - 12 hours) and plasma corticosterone levels remain elevated for 24 hours (Malendowicz et al., 1993). In addition, the stress-evoked rise in corticosterone was absent in rats treated for 6 days with NMU-8 (Malendowicz et al. 1994a). Since corticosterone exerts both mineralocorticoid and glucocorticoid effects, regulation of



its release by NMU ligands would be expected to modulate fluid homeostasis, ionic balance and metabolism. Although the mechanisms that mediate these effects remain unclear, identification of NMU-like immunoreactivity in nerve fibers in the rat hypothalamic paraventricular and supraoptic nuclei suggest a potential role for NMU in the hypothalamic regulation of pituitary function (Steel et al. 1988).

The corticosterone releasing effects of NMU may be mediated in part by direct effects on the adrenal gland. In rat adrenal gland slices, NMU-8 markedly increased basal corticosterone and pregnenolone steroid secretion (Malendowicz et al. 1994a and 1994b). These effects require the presence of adrenal medulla suggesting that NMU-8 acts on medullary chromaffin cells which may stimulate cells of the cortex through a paracrine mechanism. On the other hand, rat NMU-23 directly decreased basal corticosterone secretion from isolated rat inner adrenocortical cells (in the absence of medullary cells) while NMU-8 was without effect (Malendowicz and Nussdorfer 1993). This discrepancy between NMU-23 and NMU-8 effects on adrenal cortical cells suggests that the NMU receptor in these cells differs from that responsible for smooth muscle contraction. Repeated NMU-8 administration also decreased adrenal weight and the number of cells in the zona reticularis, further suggesting a stimulatory role for NMU on adrenal gland (Malendowicz et al. 1994a). NMU ligands may therefore be useful for directly regulating secretion from the adrenal gland.

Although NMU-like immunoreactivity has not been demonstrated within the adrenal gland or circulating in plasma, corticotrophs within the anterior lobe of rat and human pituitary gland contain high levels of NMU-like immunoreactivity (Steel et al. 1988) suggesting a possible hormonal role for NMU. Co-release of NMU with other

bioactive peptides is likely to occur since NMU was observed by electron microscopy to be present in the same secretory granules as ACTH and galanin (Cimini et al. 1993).

Furthermore, both ACTH and NMU are present in human  
5 pituitary corticotropinomas as well as in ACTH expressing tumors from a variety of other tissue sources (Steel et al. 1988). Supporting a potential hormonal role of NMU, is the identification of a small population of NMU positive parafollicular C-cells in rat thyroid gland (Domin et al. 1990 and Lo et al. 1992).

Activities of this peptide also include a hypertensive effect when given intravenously to rats at a high dose (1 nmole; Gardiner et al. 1990). However, at a lower dose (0.1 nmole), NMU caused potent constrictor effects on the superior mesenteric vascular bed reducing mesenteric blood flow without changing systemic blood pressure. The NMU-induced reduction in mesenteric blood flow was also demonstrated in dog (Sumi et al. 1987). In addition, a  
20 slight increase in blood flow to the pancreas was measured in these experiments. Such actions suggest the involvement of NMU in the regulation of blood flow to the digestive tract and subsequent effects on digestion.

NMU was originally isolated based on its potent uterine contractile activity *in vitro* and has contractile activity on other smooth muscle preparations including chicken crop (Minamino, N. et al. 1985a, 1985b). Isolated muscle strips from the dome of the human urinary bladder were also  
30 contracted by NMU (Maggi et al. 1990) suggesting a role for this peptide in urinary control. NMU-like immunoreactivity has been identified in high levels in the rat genito-urinary systems including vas deferens, prostate, fallopian tube, urethra, vagina, ovary and uterus (reviewed in Nandha and  
35 Bloom 1993). Smooth muscle contractile or other hormonal

effects of NMU in these tissues may regulate urinary control and/or reproductive functions.

Along with many other neuropeptides, NMU is present in nerves throughout the gastrointestinal tract (reviewed in Nandha and Bloom 1993). NMU stimulates contraction of isolated longitudinal muscle of human ileum (Maggi et al. 1990) and rat stomach circular muscle (Benito-Orfila et al. 1991) suggesting a role for NMU in gastric emptying and intestinal motility. Interestingly, porcine jejunum (Brown and Quito 1988) and guinea pig small intestine (Minamino et al. 1985b) are not contracted by NMU indicating species differences in gut regulation by this peptide. However, ion transport is modulated by NMU-8 in isolated porcine jejunal mucosa (Brown and Quito 1988). NMU-like immunoreactivity in the intestine has been localized to both the submucosal and myenteric ganglion cells (Ballesta et al. 1988) consistent with the observed effects on contractility, blood flow and absorptive/secretory functions (Ballesta et al. 1988).

Although higher concentrations of NMU are found in the periphery than in the central nervous system (CNS), immunocytochemical analysis demonstrated the presence of NMU in nerve fibers in many CNS regions with concentrations in discrete functional systems (Honzawa et al. 1987, Ballesta et al. 1988 and reviewed in Domin et al. 1987). For example, NMU-like immunoreactivity was identified in all of the cranial nuclei associated with somato-motor function (Ballesta et al. 1988). Several structures associated with sensory processing are also rich in NMU containing fibers including spinal cord (dorsal horn > ventral horn), trigeminal sensory nuclei, vestibular nuclei and other nuclei associated with descending spinal pathways (Honzawa et al. 1987). This localization suggests a role for NMU in perception and processing of sensory stimuli including pain.

Three cerebellar nuclei (nucleus medialis, interpositus and lateralis) also demonstrated NMU-like immunoreactivity, consistent with the potential importance of NMU in sensory processing. Neuronal cell bodies containing NMU-like immunoreactivity have been identified in the arcuate nucleus of the hypothalamus, an area identified as important for the regulation of food intake and neuroendocrine control. Relatively high levels of NMU-like immunoreactivity were also detected in the nucleus accumbens (Domin et al. 1987), an area where dopaminergic transmission is involved in reward and reinforcement of learned behaviors. The presence of NMU in another area important in dopaminergic transmission, the substantia nigra, (Domin et al. 1987) suggests a role for NMU in the modulation of dopaminergic actions in movement control as well. NMU-like immunoreactivity is also found in the hippocampus, amygdala and other portions of the limbic system suggesting a role for NMU ligands in affective disorders, psychosis and cognition.

G-protein coupled receptors (GPCR's) activated by this peptide and related analogues were postulated to exist based on binding of [<sup>125</sup>I]rat NMU in rat uterus membranes (Nandha K.A. et al. 1993). The binding is saturable and of high affinity (Kd = 0.35 nM, maximal binding capacity (Bmax) = 580 fmol/mg protein). This affinity corresponds to the EC50 of contractile activity in this tissue, 0.2 nM, consistent with the involvement of this binding site in NMU-induced uterine contraction. The GTP analogue, GTPγS, inhibited binding of [<sup>125</sup>I]rat NMU-23 suggesting that the binding site is a GPCR. In addition, chemical cross-linking identified the binding protein as having an apparent Mr of 48,500 which is consistent with the expected size of a GPCR protein. Nandha et al. (1994) also identified [<sup>125</sup>I]rat NMU-23 binding sites in rat uterus tissue slices and in the indusium

griseum by autoradiography.

09609445 063000

SUMMARY OF THE INVENTION

This invention provides an isolated nucleic acid encoding a mammalian SNORF62 receptor.

This invention also provides an isolated nucleic acid encoding a mammalian SNORF72 receptor.

This invention further provides a purified mammalian SNORF62 receptor protein.

This invention still further provides a purified mammalian SNORF72 receptor protein.

Furthermore, this invention provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF62 receptor, wherein the probe has a sequence complementary to a unique sequence present within one of the two strands of the nucleic acid encoding the human SNORF62 receptor contained in plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042).

This invention further provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF72 receptor, wherein the probe has a sequence complementary to a unique sequence present within one of the two strands of the nucleic acid encoding the human SNORF72 receptor contained in plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446).

This invention further provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF72

receptor, wherein the probe has a sequence complementary to a unique sequence present within one of the two strands of the nucleic acid encoding the rat SNORF72 receptor contained in plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927).

This invention provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF62 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 1A-1B (SEQ ID NO: 1) or (b) the reverse complement thereof.

This invention also provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF72 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 3A-3B (SEQ ID NO: 3) or (b) the reverse complement thereof.

This invention also provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF72 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 15A-15B (SEQ ID NO: 25) or (b) the reverse complement thereof.

This invention further provides a transgenic, nonhuman mammal comprising a homologous recombination knockout of the native mammalian SNORF62 receptor.

This invention still further provides a transgenic, nonhuman

mammal comprising a homologous recombination knockout of the native mammalian SNORF72 receptor.

5 This invention additionally provides a process for identifying a chemical compound which specifically binds to a mammalian SNORF62 receptor which comprises contacting cells containing DNA encoding, and expressing on their cell surface, the mammalian SNORF62 receptor, wherein such cells do not normally express the mammalian SNORF62 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF62 receptor.

10  
15  
20 Furthermore, this invention provides a process for identifying a chemical compound which specifically binds to a mammalian SNORF62 receptor which comprises contacting a membrane preparation from cells containing DNA encoding, and expressing on their cell surface, the mammalian SNORF62 receptor, wherein such cells do not normally express the mammalian SNORF62 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF62 receptor.

25 This invention further provides a process for identifying a chemical compound which specifically binds to a mammalian SNORF72 receptor which comprises contacting cells containing DNA encoding, and expressing on their cell surface, the mammalian SNORF72 receptor, wherein such cells do not normally express the mammalian SNORF72 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF72 receptor.

35 This invention provides a process for identifying a chemical



5

20

25

30

35

detecting specific binding of the chemical compound to the  
mammalian NMU receptor, a decrease in the binding of the  
second chemical compound to the mammalian NMU receptor in  
the presence of the chemical compound being tested  
5 indicating that such chemical compound binds to the  
mammalian NMU receptor.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225  
222

bind specifically to the mammalian NMU receptor under conditions permitting binding of compounds known to bind to the mammalian NMU receptor; (b) determining whether the binding of a compound known to bind to the mammalian NMU receptor is reduced in the presence of the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so (c) separately determining the binding to the mammalian NMU receptor of each compound included in the plurality of compounds, so as to thereby identify any compound included therein which specifically binds to the mammalian NMU receptor.

Furthermore, this invention provides a method for diagnosing a predisposition to a disorder associated with the activity of a specific mammalian allele which comprises: (a) obtaining DNA of subjects suffering from the disorder; (b) performing a restriction digest of the DNA with a panel of restriction enzymes; (c) electrophoretically separating the resulting DNA fragments on a sizing gel; (d) contacting the resulting gel with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a mammalian SNORF62 receptor and labeled with a detectable marker; (e) detecting labeled bands which have hybridized to the DNA encoding a mammalian SNORF62 receptor of claim 1 to create a unique band pattern specific to the DNA of subjects suffering from the disorder; (f) repeating steps (a)-(e) with DNA obtained for diagnosis from subjects not yet suffering from the disorder; and (g) comparing the unique band pattern specific to the DNA of subjects suffering from the disorder from step (e) with the band pattern from step (f) for subjects not yet suffering from the disorder so as to determine whether the patterns are the same or different and thereby diagnose predisposition to the disorder if the

patterns are the same.

09509146-063000  
This invention provides a method for diagnosing a predisposition to a disorder associated with the activity of a specific mammalian allele which comprises: (a) obtaining DNA of subjects suffering from the disorder; (b) performing a restriction digest of the DNA with a panel of restriction enzymes; (c) electrophoretically separating the resulting DNA fragments on a sizing gel; (d) contacting the resulting gel with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a mammalian SNORF72 receptor and labeled with a detectable marker; (e) detecting labeled bands which have hybridized to the DNA encoding a mammalian SNORF72 receptor of claim 2 to create a unique band pattern specific to the DNA of subjects suffering from the disorder; (f) repeating steps (a)-(e) with DNA obtained for diagnosis from subjects not yet suffering from the disorder; and (g) comparing the unique band pattern specific to the DNA of subjects suffering from the disorder from step (e) with the band pattern from step (f) for subjects not yet suffering from the disorder so as to determine whether the patterns are the same or different and thereby diagnose predisposition to the disorder if the patterns are the same.

This invention also provides a process for determining whether a chemical compound is a mammalian SNORF62 receptor agonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF62 receptor with the compound under conditions permitting the activation of the mammalian SNORF62 receptor, and detecting any increase in mammalian SNORF62 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF62 receptor agonist.

5

20

25

35

contacting cells producing a second messenger response and  
expressing on their cell surface the mammalian SNORF62  
receptor, wherein such cells do not normally express the  
mammalian SNORF62 receptor, with the chemical compound under  
conditions suitable for activation of the mammalian SNORF62  
receptor, and measuring the second messenger response in the  
presence and in the absence of the chemical compound, a  
change in the second messenger response in the presence of  
the chemical compound indicating that the compound activates  
the mammalian SNORF62 receptor.

This invention also provides a process for determining  
whether a chemical compound specifically binds to and  
activates a mammalian SNORF72 receptor, which comprises  
contacting cells producing a second messenger response and  
expressing on their cell surface the mammalian SNORF72  
receptor, wherein such cells do not normally express the  
mammalian SNORF72 receptor, with the chemical compound under  
conditions suitable for activation of the mammalian SNORF72  
receptor, and measuring the second messenger response in the  
presence and in the absence of the chemical compound, a  
change in the second messenger response in the presence of  
the chemical compound indicating that the compound activates  
the mammalian SNORF72 receptor.

This invention further provides a process for determining  
whether a chemical compound specifically binds to and  
inhibits activation of a mammalian NMU receptor, which  
comprises separately contacting cells producing a second  
messenger response and expressing on their cell surface the  
mammalian NMU receptor, wherein such cells do not normally  
express the mammalian NMU receptor, with both the chemical  
compound and a second chemical compound known to activate  
the mammalian NMU receptor, and with only the second  
chemical compound, under conditions suitable for activation

of the mammalian NMU receptor, and measuring the second messenger response in the presence of only the second chemical compound and in the presence of both the second chemical compound and the chemical compound, a smaller change in the second messenger response in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound indicating that the chemical compound inhibits activation of the mammalian NMU receptor.

This invention provides a method of screening a plurality of chemical compounds not known to activate a mammalian SNORF62 receptor to identify a compound which activates the mammalian SNORF62 receptor which comprises: (a) contacting cells transfected with and expressing the mammalian SNORF62 receptor with the plurality of compounds not known to activate the mammalian SNORF62 receptor, under conditions permitting activation of the mammalian SNORF62 receptor; (b) determining whether the activity of the mammalian SNORF62 receptor is increased in the presence of one or more of the compounds; and if so (c) separately determining whether the activation of the mammalian SNORF62 receptor is increased by any compound included in the plurality of compounds, so as to thereby identify each compound which activates the mammalian SNORF62 receptor.

This invention also provides a method of screening a plurality of chemical compounds not known to activate a mammalian SNORF72 receptor to identify a compound which activates the mammalian SNORF72 receptor which comprises: (a) contacting cells transfected with and expressing the mammalian SNORF72 receptor with the plurality of compounds not known to activate the mammalian SNORF72 receptor, under conditions permitting activation of the mammalian SNORF72 receptor; (b) determining whether the activity of the

mammalian SNORF72 receptor is increased in the presence of one or more of the compounds; and if so (c) separately determining whether the activation of the mammalian SNORF72 receptor is increased by any compound included in the plurality of compounds, so as to thereby identify each compound which activates the mammalian SNORF72 receptor.

000009445-063000  
This invention further provides a method of screening a plurality of chemical compounds not known to inhibit the activation of a mammalian NMU receptor to identify a compound which inhibits the activation of the mammalian NMU receptor, which comprises: (a) contacting cells transfected with and expressing the mammalian NMU receptor with the plurality of compounds in the presence of a known mammalian NMU receptor agonist, under conditions permitting activation of the mammalian NMU receptor; (b) determining whether the extent or amount of activation of the mammalian NMU receptor is reduced in the presence of one or more of the compounds, relative to the extent or amount of activation of the mammalian NMU receptor in the absence of such one or more compounds; and if so (c) separately determining whether each such compound inhibits activation of the mammalian NMU receptor for each compound included in the plurality of compounds, so as to thereby identify any compound included in such plurality of compounds which inhibits the activation of the mammalian NMU receptor.

This invention still further provides a method of screening a plurality of chemical compounds not known to inhibit the activation of a mammalian SNORF72 receptor to identify a compound which inhibits the activation of the mammalian SNORF72 receptor, which comprises: (a) contacting cells transfected with and expressing the mammalian SNORF72 receptor with the plurality of compounds in the presence of a known mammalian SNORF72 receptor agonist, under conditions



0909146 163000  
5 permitting activation of the mammalian SNORF72 receptor; (b) determining whether the extent or amount of activation of the mammalian SNORF72 receptor is reduced in the presence of one or more of the compounds, relative to the extent or amount of activation of the mammalian SNORF72 receptor in the absence of such one or more compounds; and if so (c) separately determining whether each such compound inhibits activation of the mammalian SNORF72 receptor for each compound included in the plurality of compounds, so as to thereby identify any compound included in such plurality of compounds which inhibits the activation of the mammalian SNORF72 receptor.

20 This invention additionally provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF62 receptor which comprises administering to the subject a compound which is a mammalian SNORF62 receptor agonist in an amount effective to treat the abnormality.

25 This invention also provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF72 receptor which comprises administering to the subject a compound which is a mammalian SNORF72 receptor agonist in an amount effective to treat the abnormality.

30 This invention further provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF62 receptor which comprises administering to the subject a compound which is a mammalian SNORF62 receptor antagonist in an amount effective to treat the abnormality.

35 This invention still further provides a method of treating

an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF72 receptor which comprises administering to the subject a compound which is a mammalian SNORF72 receptor antagonist in an amount effective to treat the abnormality.

5

00000000 063000 0909145 00000000

**BRIEF DESCRIPTION OF THE FIGURES**

**Figures 1A-1B**

5 Nucleotide sequence including sequence encoding a human SNORF62 receptor (SEQ ID NO: 1). Putative open reading frames including the longest and shortest open reading frames are indicated by underlining two start (ATG) codons (at positions 38-40 and 107-109) and the stop codon (at positions 1316-1318). In addition, partial 5' and 3' untranslated sequences are shown.

**Figures 2A-2B**

5 Deduced amino acid sequence (SEQ ID NO: 2) of the human SNORF62 receptor encoded by the longest open reading frame indicated in the nucleotide sequence shown in Figures 1A-1B (SEQ ID NO: 1). The seven putative transmembrane (TM) regions are underlined.

**Figures 3A-3B**

20 Nucleotide sequence including sequence encoding a human SNORF72 receptor (SEQ ID NO: 3). Putative open reading frames including the longest and shortest open reading frames are indicated by underlining two start (ATG) codons (at positions 27-29 and 36-38) and the stop codon (at positions 1272-1274). In addition, partial 5' and 3' untranslated sequences are shown.

**Figures 4A-4B**

30 Deduced amino acid sequence (SEQ ID NO: 4) of the human SNORF72 receptor encoded by the longest open reading frame indicated in the nucleotide sequence shown in Figures 3A-3B (SEQ ID NO: 3). The seven putative transmembrane (TM) regions are underlined.

**Figure 5**

Pairwise GAP comparison (Wisconsin Package, Genetics Computer Group, Madison, WI) of the amino acid sequences of SNORF62 (upper sequence) and SNORF72 (lower sequence). (|) Indicates identical residues, and (:) or (.) indicate varying degrees of conservation between residues.

**Figure 6**

Concentration-dependent stimulation of intracellular  $\text{Ca}^{2+}$  release by human NMU-25 in DNA vector (Mock)- and SNORF62-transfected COS-7 cells. The data presented are representative of 7 experiments performed in duplicate.

**Figure 7**

Concentration-dependent stimulation of intracellular  $\text{Ca}^{2+}$  release by human NMU-25 in SNORF72-transfected COS-7 cells. The data presented are representative of 2 experiments performed in duplicate.

**Figure 8**

Stimulation of intracellular  $\text{Ca}^{2+}$  release by NMU and related peptides (300 nM) in SNORF72-transfected COS-7 cells. The data presented are representative of 2 experiments performed in duplicate.

**Figures 9A and 9B**

Saturation binding of [ $^{125}\text{I}$ ]rat NMU-23 and [ $^{125}\text{I}$ ]NMU-8 to SNORF62. COS-7 cells were transiently transfected with SNORF62 and membranes were prepared as described in Materials and Methods. Membranes (5 - 20  $\mu\text{g}$  protein) were incubated at 4° C with increasing concentrations of [ $^{125}\text{I}$ ]rat NMU-23 (Figure 9A) or [ $^{125}\text{I}$ ]NMU-8 (Figure 9B) ( 0.01 - 3 nM) for 60 minutes. Non-specific binding was determined in the presence of 100 nM rat NMU-23. Results are representative of

2 experiments performed in duplicate.

**Figure 10**

Displacement of [ $^{125}$ I]-rat NMU-23 binding in SNORF62-transfected COS-7 membranes. Membranes were incubated with [ $^{125}$ I]-rat NMU-23 (0.05 - 0.1 nM) in the presence of the indicated peptides as described in Materials and Methods. Results presented are representative of 2 experiments.

**Figures 11A and 11B**

Saturation binding of [ $^{125}$ I]rat NMU-23 and [ $^{125}$ I]NMU-8 to SNORF72. COS-7 cells were transiently transfected with SNORF72 and membranes were prepared as described in Materials and Methods. Membranes (5 - 20  $\mu$ g protein) were incubated at 4° C with increasing concentrations of [ $^{125}$ I]rat NMU-23 (Figure 11A) or [ $^{125}$ I]NMU-8 (Figure 11B) (0.01 - 2.8 nM) for 60 minutes. Non-specific binding was determined in the presence of 100 nM rat NMU-23. Results presented are representative of 2 experiments performed in duplicate.

**Figure 12**

Displacement of [ $^{125}$ I]rat NMU-23 binding in SNORF72-transfected COS-7 membranes. Membranes were incubated with [ $^{125}$ I]rat NMU-23 (0.05 - 0.1 nM) in the presence of the indicated peptides as described in Materials and Methods. Results are representative of 2 experiments performed in duplicate.

**Figure 13**

Representative traces of human NMU-25-induced calcium-activated chloride currents in *Xenopus laevis* oocytes. The traces labeled SNORF62 were recorded from oocytes injected with mRNA encoding SNORF62.

**Figures 14A-14B**

Nucleotide sequence including sequence encoding a rat SNORF72 receptor (SEQ ID NO: 24). Putative open reading frames including the longest and shortest open reading frames are indicated by underlining two start (ATG) codons (at positions 23-25 and 65-67) and the stop codon (at positions 1208-1210). In addition, partial 5' and 3' untranslated sequences are shown.

**Figures 15A-15B**

Deduced amino acid sequence (SEQ ID NO: 25) of the rat SNORF72 receptor encoded by the longest open reading frame indicated in the nucleotide sequence shown in Figures 14A-14B (SEQ ID NO: 24). The seven putative transmembrane (TM) regions are underlined.

**Figures 16A-16B**

Amino acid sequence comparison of rat SNORF72 (SEQ ID NO: 25) with human SNORF72 (SEQ ID NO: 4) and human SNORF62 (SEQ ID NO: 2). The multiple sequence alignment was generated using Pileup (Wisconsin Package Version 10.0, Genetics Computer Group (GCG), Madison, Wisc.).

**Figures 17A and 17B**

Nucleotide sequence including sequence encoding a rat SNORF62a receptor (SEQ ID NO: 26). Putative open reading frames are indicated by underlining the start (ATG) codon (at positions 26-28) and the stop codon (at positions 1265-1267). In addition, partial 5' and 3' untranslated sequences are shown.

**Figures 18A and 18B**

Deduced amino acid sequence (SEQ ID NO: 27) of the rat SNORF62a receptor encoded by the longest open reading frame

indicated in the nucleotide sequence shown in Figures 17A-17B (SEQ ID NO: 26). The seven putative transmembrane (TM) regions are underlined.

5 **Figures 19A and 19B**

Nucleotide sequence including sequence encoding a rat SNORF62b receptor (SEQ ID NO: 28). Putative open reading frames including the longest and shortest open reading frames are indicated by underlining two start (ATG) codons (at positions 27-29 and 69-71) and the stop codon (at position 1344-1346). In addition, partial 5' and 3' untranslated sequences are shown.

10 **Figures 20A and 20B**

Deduced amino acid sequence of the rat SNORF62b receptor (SEQ ID NO: 29) encoded by the longest open reading frame indicated in the nucleotide sequence shown in Figures 19A-19B. The seven putative transmembrane (TM) regions are underlined.

15 **Figures 21A-21C**

Amino acid sequence comparison of rat SNORF62a (SEQ ID NO: 27), rat SNORF62b (SEQ ID NO: 29) and human SNORF62 (SEQ ID NO: 2). The multiple sequence alignment was generated using Pileup (Wisconsin Package Version 10.0, Genetics Computer Group (GCG), Madison, Wisc.).

20 **Figure 22**

Stimulation of intracellular  $\text{Ca}^{2+}$  release by NMU-8 and rat NMU-23 at 100 nM in rat SNORF72-transfected COS-7 cells. The data represent the average  $\pm$  SEM for an experiment performed in quadruplicate.

Concentration-dependent stimulation of inositol phosphate (IP) second messenger release by human NMU-25 in SNORF62-transfected and mock-transfected Cos-7 cells. The data presented are representative of 3 experiments performed in triplicate.

5



**DETAILED DESCRIPTION OF THE INVENTION**

000009146-063000  
5 This invention provides a recombinant nucleic acid comprising a nucleic acid encoding a mammalian SNORF62 receptor, wherein the mammalian receptor-encoding nucleic acid hybridizes under high stringency conditions to a nucleic acid encoding a human SNORF62 receptor and having a sequence identical to the sequence of the human SNORF62 receptor-encoding nucleic acid contained in plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042).

5 This invention further provides a recombinant nucleic acid comprising a nucleic acid encoding a human SNORF62 receptor, wherein the human SNORF62 receptor comprises an amino acid sequence identical to the sequence of the human SNORF62 receptor encoded by the longest open reading frame indicated in Figures 1A-1B (SEQ ID NO: 1).

20 The plasmid pEXJ.T3T7-hSNORF62-f was deposited on December 8, 1999, with the American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, Virginia 20110-2209, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of  
25 Microorganisms for the Purposes of Patent Procedure and was accorded Patent Deposit Designation No. PTA-1042.

30 This invention further provides a recombinant nucleic acid comprising a nucleic acid encoding a mammalian SNORF72 receptor, wherein the mammalian receptor-encoding nucleic acid hybridizes under high stringency conditions to a nucleic acid encoding a human SNORF72 receptor and having a sequence identical to the sequence of the human SNORF72 receptor-encoding nucleic acid contained in plasmid  
35 pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-

1446).

This invention further provides a recombinant nucleic acid comprising a nucleic acid encoding a human SNORF72 receptor, wherein the human SNORF72 receptor comprises an amino acid sequence identical to the sequence of the human SNORF72 receptor encoded by the longest open reading frame indicated in Figures 3A-3B (SEQ ID NO: 3).

The plasmid pEXJ.T3T7-hSNORF72-f was deposited on March 2, 2000, with the American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, Virginia 20110-2209, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and was accorded Patent Deposit Designation No. PTA-1446.

The plasmid pEXJ.BS-rSNORF72-f was deposited on May 26, 2000, with the American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, Virginia 20110-2209, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and was accorded Patent Deposit Designation No. PTA-1927.

This invention also contemplates recombinant nucleic acids which comprise nucleic acids encoding naturally occurring allelic variants of the mammalian SNORF62 and mammalian SNORF72 receptors described above.

Hybridization methods are well known to those of skill in the art. For purposes of this invention, hybridization under high stringency conditions means hybridization performed at 40°C in a hybridization buffer containing 50% formamide, 5X SSC, 7mM Tris, 1X Denhardt's, 25µg/ml salmon sperm DNA; wash

at 50°C in 0.1X SSC, 0.1%SDS.

Throughout this application, the following standard abbreviations are used to indicate specific nucleotide bases:

A = adenine  
G = guanine  
C = cytosine  
T = thymine  
M = adenine or cytosine  
R = adenine or guanine  
W = adenine or thymine  
S = cytosine or guanine  
Y = cytosine or thymine  
K = guanine or thymine  
V = adenine, cytosine, or guanine (not thymine)  
H = adenine, cytosine, or thymine (not cytosine)  
B = cytosine, guanine, or thymine (not adenine)  
N = adenine, cytosine, guanine, or thymine (or other modified base such as inosine)  
I = inosine

Furthermore, the term "agonist" is used throughout this application to indicate any peptide or non-peptidyl compound which increases the activity of any of the polypeptides of the subject invention. The term "antagonist" is used throughout this application to indicate any peptide or non-peptidyl compound which decreases the activity of any of the polypeptides of the subject invention.

Furthermore, as used herein, the phrase "pharmaceutically acceptable carrier" means any of the standard pharmaceutically acceptable carriers. Examples include, but are not limited to, phosphate buffered saline, physiological saline, water, and emulsions, such as oil/water emulsions.

It is possible that the mammalian SNORF62 receptor gene and the mammalian SNORF72 receptor gene contain introns and furthermore, the possibility exists that additional introns could exist in coding or non-coding regions. In addition, spliced form(s) of mRNA may encode additional amino acids either upstream of the currently defined starting methionine or within the coding region. Further, the existence and use of alternative exons is possible, whereby the mRNA may encode different amino acids within the region comprising the exon. In addition, single amino acid substitutions may arise via the mechanism of RNA editing such that the amino acid sequence of the expressed protein is different than that encoded by the original gene. (Burns, et al., 1996; Chu, et al., 1996). Such variants may exhibit pharmacologic properties differing from the polypeptide encoded by the original gene.

This invention provides splice variants of the mammalian SNORF62 and SNORF72 receptors disclosed herein. This invention further provides for alternate translation initiation sites and alternately spliced or edited variants of nucleic acids encoding the mammalian SNORF62 and SNORF72 receptors of this invention.

The nucleic acids of the subject invention also include nucleic acid analogs of the human SNORF62 receptor gene, wherein the human SNORF62 receptor gene comprises the nucleic acid sequence shown in Figures 1A-1B (SEQ ID NO: 1) or contained in plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042). Nucleic acid analogs of the human SNORF62 receptor genes differ from the human SNORF62 receptor genes described herein in terms of the identity or location of one or more nucleic acid bases (deletion analogs containing less than all of the nucleic acid bases shown in Figures 1A-1B or contained in plasmid pEXJ.T3T7-hSNORF62-f,

substitution analogs wherein one or more nucleic acid bases shown in Figures 1A-1B or contained in plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042), are replaced by other nucleic acid bases, and addition analogs, wherein one or more nucleic acid bases are added to a terminal or medial portion of the nucleic acid sequence) and which encode proteins which share some or all of the properties of the proteins encoded by the nucleic acid sequences shown in Figures 1A-1B or contained in plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042). In one embodiment of the present invention, the nucleic acid analog encodes a protein which has an amino acid sequence identical to that shown in Figures 2A-2B or encoded by the nucleic acid sequence contained in plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042). In another embodiment, the nucleic acid analog encodes a protein having an amino acid sequence which differs from the amino acid sequences shown in Figures 2A-2B or encoded by the nucleic acid contained in plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042). In a further embodiment, the protein encoded by the nucleic acid analog has a function which is the same as the function of the receptor proteins having the amino acid sequence shown in Figures 2A-2B. In another embodiment, the function of the protein encoded by the nucleic acid analog differs from the function of the receptor protein having the amino acid sequence shown in Figures 2A-2B. In another embodiment, the variation in the nucleic acid sequence occurs within the transmembrane (TM) region of the protein. In a further embodiment, the variation in the nucleic acid sequence occurs outside of the TM region.

The nucleic acids of the subject invention also include nucleic acid analogs of the rat SNORF62a and rat SNORF62b receptor genes, wherein the rat SNORF62a receptor gene

comprises the nucleic acid sequence shown in Figures 17A-17B (SEQ ID NO: 26) and the rat SNORF62b receptor gene comprises the nucleic acid sequence shown in Figures 19A-19B (SEQ ID NO: 28). Nucleic acid analogs of the rat SNORF62a and rat SNORF62b receptor genes differ from the rat SNORF62a and rat SNORF62b receptor genes described herein in terms of the identity or location of one or more nucleic acid bases (deletion analogs containing less than all of the nucleic acid bases shown in Figures 17A-17B or Figures 19A-19B, substitution analogs wherein one or more nucleic acid bases shown in Figures 17A-17B or Figures 19A-19B, are replaced by other nucleic acid bases, and addition analogs, wherein one or more nucleic acid bases are added to a terminal or medial portion of the nucleic acid sequence) and which encode proteins which share some or all of the properties of the proteins encoded by the nucleic acid sequences shown in Figures 17A-17B or Figure 19A-19B. In one embodiment of the present invention, the nucleic acid analog encodes a protein which has an amino acid sequence identical to that shown in Figures 18A-18B or Figures 20A-20B. In another embodiment, the nucleic acid analog encodes a protein having an amino acid sequence which differs from the amino acid sequences shown in Figures 18A-18B or Figures 20A-20B. In a further embodiment, the protein encoded by the nucleic acid analog has a function which is the same as the function of the receptor proteins having the amino acid sequence shown in Figures 18A-18B or Figures 20A-20B. In another embodiment, the function of the protein encoded by the nucleic acid analog differs from the function of the receptor protein having the amino acid sequence shown in Figures 18A-18B or Figures 20A-20B. In another embodiment, the variation in the nucleic acid sequence occurs within the transmembrane (TM) region of the protein. In a further embodiment, the variation in the nucleic acid sequence occurs outside of the TM region.

The nucleic acids of the subject invention also include nucleic acid analogs of the human SNORF72 receptor gene, wherein the human SNORF72 receptor gene comprises the nucleic acid sequence shown in Figures 3A-3B (SEQ ID NO: 3) or contained in plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446). Nucleic acid analogs of the human SNORF72 receptor gene differ from the human SNORF72 receptor gene described herein in terms of the identity or location of one or more nucleic acid bases (deletion analogs containing less than all of the nucleic acid bases shown in Figures 3A-3B or contained in plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446), substitution analogs wherein one or more nucleic acid bases shown in Figures 3A-3B or contained in plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446), are replaced by other nucleic acid bases, and addition analogs, wherein one or more nucleic acid bases are added to a terminal or medial portion of the nucleic acid sequence) and which encode proteins which share some or all of the properties of the proteins encoded by the nucleic acid sequences shown in Figures 3A-3B or contained in plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446). In one embodiment of the present invention, the nucleic acid analog encodes a protein which has an amino acid sequence identical to that shown in Figures 4A-4B or encoded by the nucleic acid sequence contained in plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446). In another embodiment, the nucleic acid analog encodes a protein having an amino acid sequence which differs from the amino acid sequences shown in Figures 4A-4B or encoded by the nucleic acid contained in plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446). In a further embodiment, the protein encoded by the nucleic acid analog has a function which is the same as the function of the receptor

proteins having the amino acid sequence shown in Figures 4A-4B. In another embodiment, the function of the protein encoded by the nucleic acid analog differs from the function of the receptor protein having the amino acid sequence shown in Figures 4A-4B. In another embodiment, the variation in the nucleic acid sequence occurs within the transmembrane (TM) region of the protein. In a further embodiment, the variation in the nucleic acid sequence occurs outside of the TM region.

The nucleic acids of the subject invention also include nucleic acid analogs of the rat SNORF72 receptor gene, wherein the rat SNORF72 receptor gene comprises the nucleic acid sequence shown in Figures 14A-14B (SEQ ID NO: 24) or contained in plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927). Nucleic acid analogs of the rat SNORF72 receptor gene differ from the rat SNORF72 receptor gene described herein in terms of the identity or location of one or more nucleic acid bases (deletion analogs containing less than all of the nucleic acid bases shown in Figures 14A-14B or contained in plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927), substitution analogs wherein one or more nucleic acid bases shown in Figures 14A-14B or contained in plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927), are replaced by other nucleic acid bases, and addition analogs, wherein one or more nucleic acid bases are added to a terminal or medial portion of the nucleic acid sequence) and which encode proteins which share some or all of the properties of the proteins encoded by the nucleic acid sequences shown in Figures 14A-14B or contained in plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927). In one embodiment of the present invention, the nucleic acid analog encodes a protein which has an amino acid sequence identical to that shown in Figures 15A-15B or encoded by the nucleic



acid sequence contained in plasmid pEXJ.BS-rSNORF72-f  
(Patent Deposit Designation No. PTA-1927). In another  
embodiment, the nucleic acid analog encodes a protein having  
an amino acid sequence which differs from the amino acid  
sequences shown in Figures 15A-15B or encoded by the nucleic  
acid contained in plasmid pEXJ.BS-rSNORF72-f (Patent Deposit  
Designation No. PTA-1927). In a further embodiment, the  
protein encoded by the nucleic acid analog has a function  
which is the same as the function of the receptor proteins  
having the amino acid sequence shown in Figures 15A-15B. In  
another embodiment, the function of the protein encoded by  
the nucleic acid analog differs from the function of the  
receptor protein having the amino acid sequence shown in  
Figures 15A-15B. In another embodiment, the variation in  
the nucleic acid sequence occurs within the transmembrane  
(TM) region of the protein. In a further embodiment, the  
variation in the nucleic acid sequence occurs outside of the  
TM region.

This invention provides the above-described isolated nucleic  
acid, wherein the nucleic acid is DNA. In an embodiment,  
the DNA is cDNA. In another embodiment, the DNA is genomic  
DNA. In still another embodiment, the nucleic acid is RNA.  
Methods for production and manipulation of nucleic acid  
molecules are well known in the art.

This invention further provides nucleic acid which is  
degenerate with respect to the DNA encoding any of the  
polypeptides described herein. In an embodiment, the  
nucleic acid comprises a nucleotide sequence which is  
degenerate with respect to the nucleotide sequence shown in  
Figures 1A-1B (SEQ ID NO: 1) or the nucleotide sequence  
contained in the plasmid pEXJ.T3T7-hSNORF62-f (Patent  
Deposit Designation No. PTA-1042), that is, a nucleotide  
sequence which is translated into the same amino acid

sequence. In another embodiment, the nucleic acid comprises a nucleotide sequence which is degenerate with respect to the nucleotide sequence shown in Figures 3A-3B (SEQ ID NO: 3) or the nucleotide sequence contained in the plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446), that is, a nucleotide sequence which is translated into the same amino acid sequence. In a further embodiment, the nucleic acid comprises a nucleotide sequence which is degenerate with respect to the nucleotide sequence shown in Figures 14A-14B (SEQ ID NO: 24) or the nucleotide sequence contained in the plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927), that is, a nucleotide sequence which is translated into the same amino acid sequence. In another embodiment, the nucleic acid comprises a nucleotide sequence which is degenerate with respect to the nucleotide sequence shown in Figures 17A-17B (SEQ ID NO: 26), that is, a nucleotide sequence which is translated into the same amino acid sequence. In yet another embodiment, the nucleic acid comprises a nucleotide sequence which is degenerate with respect to the nucleotide sequence shown in Figures 19A-19B (SEQ ID NO: 28), that is, a nucleotide sequence which is translated into the same amino acid sequence.

This invention also encompasses DNAs and cDNAs which encode amino acid sequences which differ from those of the polypeptides of this invention, but which should not produce phenotypic changes.

Alternately, this invention also encompasses DNAs, cDNAs, and RNAs which hybridize to the DNA, cDNA, and RNA of the subject invention. Hybridization methods are well known to those of skill in the art.

The nucleic acids of the subject invention also include

5  
0  
15  
20

25

30

35

large scale synthesis of the polypeptides by a variety of recombinant techniques. The nucleic acid molecule is useful for generating new cloning and expression vectors, transformed and transfected prokaryotic and eukaryotic host cells, and new and useful methods for cultured growth of such host cells capable of expression of the polypeptide and related products.

This invention also provides an isolated nucleic acid encoding species homologs of the SNORF62 receptor encoded by the nucleic acid sequence shown in Figures 1A-1B (SEQ ID NO: 1) or encoded by the plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042). In one embodiment, the nucleic acid encodes a mammalian SNORF62 receptor homolog which has substantially the same amino acid sequence as does the SNORF62 receptor encoded by the plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042). In another embodiment, the nucleic acid encodes a mammalian SNORF62 receptor homolog which has above 75% amino acid identity to the SNORF62 receptor encoded by the plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042); preferably above 85% amino acid identity to the SNORF62 receptor encoded by the plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042); most preferably above 95% amino acid identity to the SNORF62 receptor encoded by the plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042). In another embodiment, the mammalian SNORF62 receptor homolog has above 70% nucleic acid identity to the SNORF62 receptor gene contained in plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042); preferably above 80% nucleic acid identity to the SNORF62 receptor gene contained in the plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042); more preferably above 90% nucleic acid identity to the SNORF62 receptor gene contained in the plasmid pEXJ.T3T7-hSNORF62-f

(Patent Deposit Designation No. PTA-1042). Examples of methods for isolating and purifying species homologs are described elsewhere (e.g., U.S. Patent No. 5,602,024, WO94/14957, WO97/26853, WO98/15570).

This invention also provides an isolated nucleic acid encoding species homologs of the SNORF62 receptor encoded by the nucleic acid sequence shown in Figures 17A-17B (SEQ ID NO: 26) or Figures 19A-19B (SEQ ID NO: 28).

This invention also provides an isolated nucleic acid encoding species homologs of the SNORF72 receptors encoded by the nucleic acid sequence shown in Figures 3A-3B (SEQ ID NO: 3) or encoded by the plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446). In one embodiment, the nucleic acid encodes a mammalian SNORF72 receptor homolog which has substantially the same amino acid sequence as does the SNORF72 receptor encoded by the plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446). In another embodiment, the nucleic acid encodes a mammalian SNORF72 receptor homolog which has above 75% amino acid identity to the SNORF72 receptor encoded by the plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446); preferably above 85% amino acid identity to the SNORF72 receptor encoded by the plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446); most preferably above 95% amino acid identity to the SNORF72 receptor encoded by the plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446). In another embodiment, the mammalian SNORF72 receptor homolog has above 70% nucleic acid identity to the SNORF72 receptor gene contained in plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446); preferably above 80% nucleic acid identity to the SNORF72 receptor gene contained in the plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446); more

preferably above 90% nucleic acid identity to the SNORF72 receptor gene contained in the plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446).

5 This invention also provides an isolated nucleic acid encoding species homologs of the SNORF72 receptors encoded by the nucleic acid sequence shown in Figures 14A-14B (SEQ ID NO: ) or encoded by the plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927). In one  
10 embodiment, the nucleic acid encodes a mammalian SNORF72 receptor homolog which has substantially the same amino acid sequence as does the SNORF72 receptor encoded by the plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927). In another embodiment, the nucleic acid encodes a  
15 mammalian SNORF72 receptor homolog which has above 75% amino acid identity to the SNORF72 receptor encoded by the plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927); preferably above 85% amino acid identity to the SNORF72 receptor encoded by the plasmid pEXJ.BS-rSNORF72-f  
20 (Patent Deposit Designation No. PTA-1927); most preferably above 95% amino acid identity to the SNORF72 receptor encoded by the plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927). In another embodiment, the mammalian SNORF72 receptor homolog has above 70% nucleic  
25 acid identity to the SNORF72 receptor gene contained in plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927); preferably above 80% nucleic acid identity to the SNORF72 receptor gene contained in the plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927); more  
30 preferably above 90% nucleic acid identity to the SNORF72 receptor gene contained in the plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927).

This invention provides an isolated nucleic acid encoding a  
35 modified mammalian SNORF62 or SNORF72 receptor, which

05609146-063000

differs from a mammalian SNORF62 or SNORF72 receptor by having an amino acid(s) deletion, replacement, or addition in the third intracellular domain.

5 This invention provides an isolated nucleic acid encoding a mammalian NMU receptor. This invention provides an isolated nucleic acid encoding a mammalian SNORF62 receptor. This invention further provides an isolated nucleic acid encoding a mammalian SNORF72 receptor. In one embodiment, the nucleic acid is DNA. In another embodiment, the DNA is cDNA. In another embodiment, the DNA is genomic DNA. In another embodiment, the nucleic acid is RNA.

10 In one embodiment, the mammalian NMU receptor is a human NMU receptor. In a further embodiment, the human NMU receptor is a human SNORF62 receptor, a rat SNORF62a receptor, or a rat SNORF62b receptor. In another embodiment, the human SNORF62 receptor has an amino acid sequence identical to that encoded by the plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042). In another embodiment, the human SNORF62 receptor has an amino acid sequence identical to the amino acid sequence shown in Figures 2A-2B (SEQ ID NO: 2). In another embodiment, the rat SNORF62a receptor has an amino acid sequence identical to the amino acid sequence shown in Figures 18A-18B (SEQ ID NO: 27). In another embodiment, the rat SNORF62b receptor has an amino acid sequence identical to the amino acid sequence shown in Figures 20A-20B (SEQ ID NO: 29).

25 In a further embodiment, the human NMU receptor is a human SNORF72 receptor or a rat SNORF72 receptor. In another embodiment, the human SNORF72 receptor has an amino acid sequence identical to that encoded by the plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446). In another embodiment, the human SNORF72 receptor has an amino

acid sequence identical to the amino acid sequence shown in  
Figures 4A-4B (SEQ ID NO: 4). In another embodiment, the  
rat SNORF72 receptor has an amino acid sequence identical to  
that encoded by the plasmid pEXJ.BS-rSNORF72-f (Patent  
Deposit Designation No. PTA-1927). In another embodiment,  
the rat SNORF72 receptor has an amino acid sequence  
identical to the amino acid sequence shown in Figures 15A-  
15B (SEQ ID NO: 25).

This invention provides a purified mammalian SNORF62 or  
SNORF72 receptor protein. In one embodiment, the SNORF62  
receptor protein is a human SNORF62 receptor protein, a rat  
SNORF62a receptor protein, or a rat SNORF62b receptor  
protein. In another embodiment, the SNORF72 receptor  
protein is a human SNORF72 receptor protein or a rat SNORF72  
receptor protein.

This invention provides a vector comprising the nucleic acid  
of this invention. This invention further provides a vector  
adapted for expression in a cell which comprises the  
regulatory elements necessary for expression of the nucleic  
acid in the cell operatively linked to the nucleic acid  
encoding the receptor so as to permit expression thereof,  
wherein the cell is a bacterial, amphibian, yeast, insect or  
mammalian cell. In one embodiment, the vector is a  
baculovirus. In another embodiment, the vector is a  
plasmid.

This invention provides a plasmid designated pEXJ.T3T7-  
hSNORF62-f (Patent Deposit Designation No. PTA-1042). This  
invention also provides a plasmid designated pEXJ.T3T7-  
hSNORF72-f (Patent Deposit Designation No. PTA-1446). This  
invention also provides a plasmid designated pEXJ.BS-  
rSNORF72-f (Patent Deposit Designation No. PTA-1927).



5

15

25

30

35

This invention further provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF62 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 1A-1B (SEQ ID NO: 1), (b) the nucleic acid sequence shown in Figures 17A-17B (SEQ ID NO: 26), (c) the nucleic acid sequence shown in Figures 19A-19B (SEQ ID NO: 28) or (d) the reverse complement to (a), (b) or (c). This invention further provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF72 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 3A-3B (SEQ ID NO: 3), (b) the nucleic acid sequence shown in Figures 14A-14B or (c) the reverse complement or (a) or (b). In one embodiment, the nucleic acid is DNA. In another embodiment, the nucleic acid is RNA.

The nucleic acids of this invention may be used as probes to obtain homologous nucleic acids from other species and to detect the existence of nucleic acids having complementary sequences in samples.

The nucleic acids may also be used to express the receptors

they encode in transfected cells.

The use of a constitutively active receptor encoded by SNORF62 either occurring naturally without further modification or after appropriate point mutations, deletions or the like, allows screening for antagonists and *in vivo* use of such antagonists to attribute a role to receptor SNORF62 without prior knowledge of the endogenous ligand.

The use of a constitutively active receptor encoded by SNORF72 either occurring naturally without further modification or after appropriate point mutations, deletions or the like, allows screening for antagonists and *in vivo* use of such antagonists to attribute a role to receptor SNORF72 without prior knowledge of the endogenous ligand.

Use of the nucleic acids further enables elucidation of possible receptor diversity and of the existence of multiple subtypes within a family of receptors of which SNORF62 is a member.

Use of the nucleic acids further enables elucidation of possible receptor diversity and of the existence of multiple subtypes within a family of receptors of which SNORF72 is a member.

Finally, it is contemplated that the receptors of this invention will serve as a valuable tool for designing drugs for treating various pathophysiological conditions such as chronic and acute inflammation, arthritis, autoimmune diseases, transplant rejection, graft vs. host disease, bacterial, fungal, protozoan and viral infections, septicemia, AIDS, pain, psychotic and neurological disorders, including anxiety, depression, schizophrenia, dementia, mental retardation, memory loss, epilepsy,

neuromotor disorders, locomotor disorders, respiratory disorders, asthma, eating/body weight disorders including obesity, bulimia, diabetes, anorexia, nausea, hypertension, hypotension, vascular and cardiovascular disorders, ischemia, stroke, cancers, ulcers, urinary retention, sexual/reproductive disorders, circadian rhythm disorders, renal disorders, bone diseases including osteoporosis, benign prostatic hypertrophy, gastrointestinal disorders, nasal congestion, dermatological disorders such as psoriasis, allergies, Parkinson's disease, Alzheimer's disease, acute heart failure, angina disorders, delirium, dyskinesias such as Huntington's disease or Gille's de la Tourette's syndrome, among others and diagnostic assays for such conditions. The receptors of this invention may also serve as a valuable tool for designing drugs for chemoprevention.

Methods of transfecting cells e.g. mammalian cells, with such nucleic acid to obtain cells in which the receptor is expressed on the surface of the cell are well known in the art. (See, for example, U.S. Patent Nos. 5,053,337; 5,155,218; 5,360,735; 5,472,866; 5,476,782; 5,516,653; 5,545,549; 5,556,753; 5,595,880; 5,602,024; 5,639,652; 5,652,113; 5,661,024; 5,766,879; 5,786,155; and 5,786,157, the disclosures of which are hereby incorporated by reference in their entireties into this application.)

Such transfected cells may also be used to test compounds and screen compound libraries to obtain compounds which bind to the SNORF62 or SNORF72 receptor, as well as compounds which activate or inhibit activation of functional responses in such cells, and therefore are likely to do so in vivo. (See, for example, U.S. Patent Nos. 5,053,337; 5,155,218; 5,360,735; 5,472,866; 5,476,782; 5,516,653; 5,545,549; 5,556,753; 5,595,880; 5,602,024; 5,639,652; 5,652,113;

5,661,024; 5,766,879; 5,786,155; and 5,786,157, the disclosures of which are hereby incorporated by reference in their entireties into this application.)

5 This invention provides an antibody capable of binding to a mammalian SNORF62 receptor encoded by a nucleic acid encoding a mammalian SNORF62 receptor. This invention further provides an antibody capable of binding to a mammalian SNORF72 receptor encoded by a nucleic acid encoding a mammalian SNORF72 receptor. In an embodiment of the present invention, the mammalian SNORF62 receptor is a human SNORF62 receptor, a rat SNORF 62a receptor, or a rat SNORF62b receptor. In a further embodiment, the mammalian SNORF72 receptor is a human SNORF72 receptor or a rat SNORF72 receptor.

10 This invention also provides an agent capable of competitively inhibiting the binding of the antibody to a mammalian SNORF62 or SNORF72 receptor. In one embodiment, the antibody is a monoclonal antibody or antisera.

15 Methods of preparing and employing antisense oligonucleotides, antibodies, nucleic acid probes and transgenic animals directed to the SNORF62 and SNORF72 receptors are well known in the art. (See, for example, U.S. Patent Nos. 5,053,337; 5,155,218; 5,360,735; 5,472,866; 5,476,782; 5,516,653; 5,545,549; 5,556,753; 5,595,880; 5,602,024; 5,639,652; 5,652,113; 5,661,024; 5,766,879; 5,786,155; and 5,786,157, the disclosures of which are hereby incorporated by reference in their entireties into this application.)

20 This invention provides for an antisense oligonucleotide having a sequence capable of specifically hybridizing to RNA encoding a mammalian SNORF62 or SNORF72 receptor, so as to

prevent translation of such RNA. This invention further provides for an antisense oligonucleotide having a sequence capable of specifically hybridizing to genomic DNA encoding a mammalian SNORF62 or SNORF72 receptor, so as to prevent transcription of such genomic DNA. In one embodiment, the oligonucleotide comprises chemically modified nucleotides or nucleotide analogues.

This invention still further provides a pharmaceutical composition comprising (a) an amount of an oligonucleotide in accordance with this invention capable of passing through a cell membrane and effective to reduce expression of a mammalian SNORF62 or SNORF72 receptor and (b) a pharmaceutically acceptable carrier capable of passing through the cell membrane.

In one embodiment, the oligonucleotide is coupled to a substance which inactivates mRNA. In another embodiment, the substance which inactivates mRNA is a ribozyme. In another embodiment, the pharmaceutically acceptable carrier comprises a structure which binds to a mammalian SNORF62 or SNORF72 receptor on a cell capable of being taken up by the cells after binding to the structure. In another embodiment, the pharmaceutically acceptable carrier is capable of binding to a mammalian SNORF62 or SNORF72 receptor which is specific for a selected cell type.

This invention also provides a pharmaceutical composition which comprises an amount of an antibody in accordance with this invention effective to block binding of a ligand to a human SNORF62 receptor or a human SNORF72 receptor and a pharmaceutically acceptable carrier.

This invention further provides a transgenic, nonhuman mammal expressing DNA encoding a mammalian SNORF62 or

SNORF72 receptor in accordance with this invention. This invention provides a transgenic, nonhuman mammal comprising a homologous recombination knockout of a native mammalian SNORF62 or SNORF72 receptor. This invention further  
5 provides a transgenic, nonhuman mammal whose genome comprises antisense DNA complementary to DNA encoding a mammalian SNORF62 or SNORF72 receptor in accordance with this invention so placed within such genome as to be transcribed into antisense mRNA which is complementary and hybridizes with mRNA encoding the mammalian SNORF62 or  
10 SNORF72 receptor so as to thereby reduce translation of such mRNA and expression of such receptor. In one embodiment, the DNA encoding the mammalian SNORF62 or SNORF72 receptor additionally comprises an inducible promoter. In another embodiment, the DNA encoding the mammalian SNORF62 or  
15 SNORF72 receptor additionally comprises tissue specific regulatory elements. In another embodiment, the transgenic, nonhuman mammal is a mouse.

20 This invention provides for a process for identifying a chemical compound which specifically binds to a mammalian SNORF62 receptor which comprises contacting cells containing DNA encoding, and expressing on their cell surface, the mammalian SNORF62 receptor, wherein such cells do not  
25 normally express the mammalian SNORF62 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF62 receptor.

30 This invention provides for a process for identifying a chemical compound which specifically binds to a mammalian SNORF72 receptor which comprises contacting cells containing DNA encoding, and expressing on their cell surface, the mammalian SNORF72 receptor, wherein such cells do not  
35 normally express the mammalian SNORF72 receptor, with the

compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF72 receptor.

5 This invention further provides for a process for identifying a chemical compound which specifically binds to a mammalian SNORF62 receptor which comprises contacting a membrane preparation from cells containing DNA encoding and expressing on their cell surface the mammalian SNORF62  
10 receptor, wherein such cells do not normally express the mammalian SNORF62 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF62 receptor.

15 This invention further provides for a process for identifying a chemical compound which specifically binds to a mammalian SNORF72 receptor which comprises contacting a membrane preparation from cells containing DNA encoding and  
20 expressing on their cell surface the mammalian SNORF72 receptor, wherein such cells do not normally express the mammalian SNORF72 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF72  
25 receptor.

In an embodiment, the mammalian SNORF62 receptor is a human SNORF62 receptor, a rat SNORF62a receptor, or a rat SNORF62b  
30 receptor. In another embodiment, the mammalian SNORF62 receptor has substantially the same amino acid sequence as the human SNORF62 receptor encoded by plasmid pEXJ.T3T7-hSNORF62-f (Patent Deposit Designation No. PTA-1042).

35 In a further embodiment, the mammalian SNORF72 receptor is a human SNORF72 receptor or a rat SNORF72 receptor. In



another embodiment, the mammalian SNORF72 receptor has substantially the same amino acid sequence as the human SNORF72 receptor encoded by plasmid pEXJ.T3T7-hSNORF72-f (Patent Deposit Designation No. PTA-1446). In yet another  
5 embodiment, the mammalian SNORF72 receptor has substantially the same amino acid sequence as the rat SNORF72 receptor encoded by plasmid pEXJ.BS-rSNORF72-f (Patent Deposit Designation No. PTA-1927).

0 In another embodiment, the mammalian SNORF62 or SNORF72 receptor has substantially the same amino acid sequence as that shown in Figures 2A-2B (SEQ ID NO: 2), Figures 4A-4B (SEQ ID NO: 4), Figures 15A-15B (SEQ ID NO: 25), Figures 18A-18B (SEQ ID NO: 27), or Figures 20A-20B (SEQ ID NO: 29),  
15 respectively. In another embodiment, the mammalian SNORF62 or SNORF72 receptor has the amino acid sequence shown in Figures 2A-2B (SEQ ID NO: 2), Figures 4A-4B (SEQ ID NO: 4), Figures 15A-15B (SEQ ID NO: 25), Figures 18A-18B (SEQ ID NO: 27), or Figures 20A-20B (SEQ ID NO: 29), respectively.

20 In one embodiment, the compound is not previously known to bind to a mammalian SNORF62 or SNORF72 receptor. In one embodiment, the cell is an insect cell. In one embodiment, the cell is a mammalian cell. In another embodiment, the  
25 cell is nonneuronal in origin. In another embodiment, the nonneuronal cell is a COS-7 cell, 293 human embryonic kidney cell, a CHO cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell. In another embodiment, the compound is a compound not previously known to bind to a mammalian SNORF62  
30 or SNORF72 receptor. This invention provides a compound identified by the preceding processes according to this invention.

35 This invention still further provides a process involving competitive binding for identifying a chemical compound

which specifically binds to a mammalian NMU receptor which comprises separately contacting cells expressing on their cell surface the mammalian NMU receptor, wherein such cells do not normally express the mammalian NMU receptor, with both the chemical compound and a second chemical compound known to bind to the receptor, and with only the second chemical compound, under conditions suitable for binding of such compounds to the receptor, and detecting specific binding of the chemical compound to the mammalian NMU receptor, a decrease in the binding of the second chemical compound to the mammalian NMU receptor in the presence of the chemical compound being tested indicating that such chemical compound binds to the mammalian NMU receptor.

This invention provides a process involving competitive binding for identifying a chemical compound which specifically binds to a mammalian NMU receptor which comprises separately contacting a membrane preparation from cells expressing on their cell surface the mammalian NMU receptor, wherein such cells do not normally express the mammalian NMU receptor, with both the chemical compound and a second chemical compound known to bind to the receptor, and with only the second chemical compound, under conditions suitable for binding of such compounds to the receptor, and detecting specific binding of the chemical compound to the mammalian NMU receptor, a decrease in the binding of the second chemical compound to the mammalian NMU receptor in the presence of the chemical compound being tested indicating that such chemical compound binds to the mammalian NMU receptor.

In an embodiment of the present invention, the second chemical compound is an NMU peptide. Examples of NMU peptides include, but are not limited to, human NMU-25, human NMU-8, porcine NMU-8, porcine NMU-25, rat NMU-25 and

any peptide comprising the carboxyl terminal seven amino acid residues of human NMU-8.

060946-063000  
5 In one embodiment, the mammalian NMU receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In another embodiment, the mammalian NMU receptor is a human SNORF72 receptor or a rat SNORF72 receptor. In a further embodiment, the cell is an insect cell. In another embodiment, the cell is a mammalian cell. In another  
10 embodiment, the cell is nonneuronal in origin. In another embodiment, the nonneuronal cell is a COS-7 cell, 293 human embryonic kidney cell, a CHO cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell. In another embodiment, the compound is not previously known to bind to a mammalian NMU  
15 receptor. This invention provides for a compound identified by the preceding process according to this invention.

20 This invention provides for a method of screening a plurality of chemical compounds not known to bind to a mammalian NMU receptor to identify a compound which specifically binds to the mammalian NMU receptor, which comprises (a) contacting cells transfected with, and expressing, DNA encoding the mammalian NMU receptor with a compound known to bind specifically to the mammalian NMU  
25 receptor; (b) contacting the cells of step (a) with the plurality of compounds not known to bind specifically to the mammalian NMU receptor, under conditions permitting binding of compounds known to bind to the mammalian NMU receptor; (c) determining whether the binding of the compound known to  
30 bind to the mammalian NMU receptor is reduced in the presence of the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so (d) separately determining the binding to the mammalian NMU receptor of each compound included in  
35 the plurality of compounds, so as to thereby identify any

compound included therein which specifically binds to the mammalian NMU receptor.

5 This invention provides a method of screening a plurality of chemical compounds not known to bind to a mammalian NMU receptor to identify a compound which specifically binds to the mammalian NMU receptor, which comprises (a) contacting a membrane preparation from cells transfected with, and expressing, DNA encoding the mammalian NMU receptor with the plurality of compounds not known to bind specifically to the mammalian NMU receptor under conditions permitting binding of compounds known to bind to the mammalian NMU receptor; (b) determining whether the binding of a compound known to bind to the mammalian NMU receptor is reduced in the presence of the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so (c) separately determining the binding to the mammalian NMU receptor of each compound included in the plurality of compounds, so as to thereby identify any compound included therein which specifically binds to the mammalian NMU receptor.

10 In one embodiment, the mammalian NMU receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In a further embodiment, the mammalian NMU receptor is a human SNORF72 receptor or a rat SNORF72 receptor. In another embodiment, the cell is a mammalian cell. In another embodiment, the mammalian cell is non-neuronal in origin. In a further embodiment, the non-neuronal cell is a COS-7 cell, a 293 human embryonic kidney cell, a LM(tk-) cell, a CHO cell, a mouse Y1 cell, or an NIH-3T3 cell.

15 This invention also provides a method of detecting expression of a mammalian SNORF62 or SNORF72 receptor by

5

[illegible]

20

30

Moreover, this invention provides method for identifying an antagonist capable of alleviating an abnormality wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF62 or SNORF72 receptor comprising  
5 administering a compound to a transgenic, nonhuman mammal according to this invention, and determining whether the compound alleviates any physiological and/or behavioral abnormality displayed by the transgenic, nonhuman mammal as a result of overactivity of a mammalian SNORF62 or SNORF72  
10 receptor, the alleviation of such an abnormality identifying the compound as an antagonist. In an embodiment, the mammalian SNORF62 receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In another embodiment, the mammalian SNORF72 receptor is a  
15 human SNORF72 receptor or a rat SNORF72 receptor.

The invention also provides an antagonist identified by the preceding method according to this invention. This invention further provides a composition, e.g. a pharmaceutical  
20 composition comprising an antagonist according to this invention and a carrier, e.g. a pharmaceutically acceptable carrier.

This invention provides a method of treating an abnormality  
25 in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF62 receptor which comprises administering to the subject an effective amount of the pharmaceutical composition according to this invention so as to thereby treat the abnormality.

This invention provides a method of treating an abnormality  
30 in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF72 receptor which comprises administering to the subject an effective  
35 amount of the pharmaceutical composition according to this

invention so as to thereby treat the abnormality.

In addition, this invention provides a method for identifying an agonist capable of alleviating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF62 or SNORF72 receptor comprising administering a compound to a transgenic, nonhuman mammal according to this invention, and determining whether the compound alleviates any physiological and/or behavioral abnormality displayed by the transgenic, nonhuman mammal, the alleviation of such an abnormality identifying the compound as an agonist. In an embodiment, the mammalian SNORF62 receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In a further embodiment, the mammalian SNORF72 receptor is a human SNORF72 receptor or a rat SNORF72 receptor. This invention provides an agonist identified by the preceding method according to this invention. This invention provides a composition, e.g. a pharmaceutical composition comprising an agonist identified by a method according to this invention and a carrier, e.g. a pharmaceutically acceptable carrier.

Moreover, this invention provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF62 or SNORF72 receptor which comprises administering to the subject an effective amount of the pharmaceutical composition of this invention so as to thereby treat the abnormality.

Yet further, this invention provides a method for diagnosing a predisposition to a disorder associated with the activity of a specific mammalian allele which comprises: (a) obtaining DNA of subjects suffering from the disorder; (b)

performing a restriction digest of the DNA with a panel of restriction enzymes; (c) electrophoretically separating the resulting DNA fragments on a sizing gel; (d) contacting the resulting gel with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a mammalian SNORF62 or SNORF72 receptor and labeled with a detectable marker; (e) detecting labeled bands which have hybridized to the DNA encoding a mammalian SNORF62 or SNORF72 receptor to create a unique band pattern specific to the DNA of subjects suffering from the disorder; (f) repeating steps (a)-(e) with DNA obtained for diagnosis from subjects not yet suffering from the disorder; and (g) comparing the unique band pattern specific to the DNA of subjects suffering from the disorder from step (e) with the band pattern from step (f) for subjects not yet suffering from the disorder so as to determine whether the patterns are the same or different and thereby diagnose predisposition to the disorder if the patterns are the same.

In one embodiment, the disorder is a disorder associated with the activity of a specific mammalian allele is diagnosed.

This invention also provides a method of preparing a purified mammalian SNORF62 receptor according to this invention which comprises: (a) culturing cells which express the mammalian SNORF62 receptor; (b) recovering the mammalian SNORF62 receptor from the cells; and (c) purifying the mammalian SNORF62 receptor so recovered.

This invention also provides a method of preparing a purified mammalian SNORF72 receptor according to this invention which comprises: (a) culturing cells which express the mammalian SNORF72 receptor; (b) recovering the mammalian



SNORF72 receptor from the cells; and (c) purifying the mammalian SNORF72 receptor so recovered.

5 This invention further provides a method of preparing a purified mammalian SNORF62 receptor according to this invention which comprises: (a) inserting a nucleic acid encoding the mammalian SNORF62 receptor into a suitable expression vector; (b) introducing the resulting vector into a suitable host cell; (c) placing the resulting host cell in  
10 suitable condition permitting the production of the mammalian SNORF62 receptor; (d) recovering the mammalian SNORF62 receptor so produced; and optionally (e) isolating and/or purifying the mammalian SNORF62 receptor so recovered.

15 This invention further provides a method of preparing a purified mammalian SNORF72 receptor according to this invention which comprises: (a) inserting a nucleic acid encoding the mammalian SNORF72 receptor into a suitable expression vector; (b) introducing the resulting vector into  
20 a suitable host cell; (c) placing the resulting host cell in suitable condition permitting the production of the mammalian SNORF72 receptor; (d) recovering the mammalian SNORF72 receptor so produced; and optionally (e) isolating  
25 and/or purifying the mammalian SNORF72 receptor so recovered.

30 Furthermore, this invention provides a process for determining whether a chemical compound is a mammalian SNORF62 receptor agonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF62 receptor with the compound under conditions  
35 permitting the activation of the mammalian SNORF62 receptor, and detecting any increase in mammalian SNORF62 receptor activity, so as to thereby determine whether the compound is

a mammalian SNORF62 receptor agonist.

Furthermore, this invention provides a process for determining whether a chemical compound is a mammalian NMU receptor agonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian NMU receptor with the compound under conditions permitting the activation of the mammalian NMU receptor, and detecting any increase in mammalian NMU receptor activity, so as to thereby determine whether the compound is a mammalian NMU receptor agonist.

Furthermore, this invention provides a process for determining whether a chemical compound is a mammalian SNORF72 receptor agonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF72 receptor with the compound under conditions permitting the activation of the mammalian SNORF72 receptor, and detecting any increase in mammalian SNORF72 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF72 receptor agonist.

This invention also provides a process for determining whether a chemical compound is a mammalian NMU receptor antagonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian NMU receptor with the compound in the presence of a known mammalian NMU receptor agonist, under conditions permitting the activation of the mammalian NMU receptor, and detecting any decrease in mammalian NMU receptor activity, so as to thereby determine whether the compound is a mammalian NMU receptor antagonist.

This invention also provides a process for determining whether a chemical compound is a mammalian SNORF62 receptor antagonist which comprises contacting cells transfected with

and expressing DNA encoding the mammalian SNORF62 receptor with the compound in the presence of a known mammalian SNORF62 receptor agonist, under conditions permitting the activation of the mammalian SNORF62 receptor, and detecting any decrease in mammalian SNORF62 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF62 receptor antagonist.

This invention also provides a process for determining whether a chemical compound is a mammalian SNORF72 receptor antagonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF72 receptor with the compound in the presence of a known mammalian SNORF72 receptor agonist, under conditions permitting the activation of the mammalian SNORF72 receptor, and detecting any decrease in mammalian SNORF72 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF72 receptor antagonist.

In an embodiment, the mammalian NMU receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In another embodiment, the mammalian NMU receptor is a human SNORF72 receptor or a rat SNORF72 receptor. In yet another embodiment, the mammalian SNORF62 receptor is a human SNORF62 receptor and the mammalian SNORF72 receptor is a human SNORF72 receptor.

This invention still further provides a composition, for example a pharmaceutical composition, which comprises an amount of a mammalian SNORF62 or SNORF72 receptor agonist determined by a process according to this invention effective to increase activity of a mammalian SNORF62 or SNORF72 receptor and a carrier, for example, a pharmaceutically acceptable carrier. In one embodiment, the mammalian SNORF62 or SNORF72 receptor agonist is not

previously known.

Also, this invention provides a composition, for example a pharmaceutical composition, which comprises an amount of a mammalian NMU receptor antagonist determined by a process according to this invention effective to reduce activity of a mammalian NMU receptor and a carrier, for example, a pharmaceutically acceptable carrier. Also, this invention provides a composition, for example a pharmaceutical composition, which comprises an amount of a mammalian SNORF62 receptor antagonist determined by a process according to this invention effective to reduce activity of a mammalian SNORF62 receptor and a carrier, for example, a pharmaceutically acceptable carrier. Also, this invention provides a composition, for example a pharmaceutical composition, which comprises an amount of a mammalian SNORF72 receptor antagonist determined by a process according to this invention effective to reduce activity of a mammalian SNORF72 receptor and a carrier, for example, a pharmaceutically acceptable carrier.

In one embodiment, the mammalian NMU receptor antagonist is not previously known. In an embodiment, the mammalian NMU receptor antagonist is a mammalian SNORF62 receptor antagonist. In a further embodiment, the mammalian NMU receptor antagonist is a mammalian SNORF72 receptor antagonist.

This invention moreover provides a process for determining whether a chemical compound specifically binds to and activates a mammalian SNORF62 receptor, which comprises contacting cells producing a second messenger response and expressing on their cell surface the mammalian SNORF62 receptor, wherein such cells do not normally express the mammalian SNORF62 receptor, with the chemical compound under

conditions suitable for activation of the mammalian SNORF62 receptor, and measuring the second messenger response in the presence and in the absence of the chemical compound, a change, e.g. an increase, in the second messenger response in the presence of the chemical compound indicating that the compound activates the mammalian SNORF62 receptor.

This invention moreover provides a process for determining whether a chemical compound specifically binds to and activates a mammalian SNORF72 receptor, which comprises contacting cells producing a second messenger response and expressing on their cell surface the mammalian SNORF72 receptor, wherein such cells do not normally express the mammalian SNORF72 receptor, with the chemical compound under conditions suitable for activation of the mammalian SNORF72 receptor, and measuring the second messenger response in the presence and in the absence of the chemical compound, a change, e.g. an increase, in the second messenger response in the presence of the chemical compound indicating that the compound activates the mammalian SNORF72 receptor.

In one embodiment, the second messenger response comprises chloride channel activation and the change in second messenger is an increase in the level of chloride current.

In another embodiment, the second messenger response comprises change in intracellular calcium levels and the change in second messenger is an increase in the measure of intracellular calcium. In another embodiment, the second messenger response comprises release of inositol phosphate and the change in second messenger is an increase in the level of inositol phosphate. In another embodiment, the second messenger response comprises release of arachidonic acid and the change in second messenger is an increase in the level of arachidonic acid. In yet another embodiment, the second messenger response comprises GTPyS ligand binding

and the change in second messenger is an increase in GTPyS ligand binding. In another embodiment, the second messenger response comprises activation of MAP kinase and the change in second messenger response is an increase in MAP kinase activation. In a further embodiment, the second messenger response comprises cAMP accumulation and the change in second messenger response is a reduction in cAMP accumulation.

This invention still further provides a process for determining whether a chemical compound specifically binds to and inhibits activation of a mammalian NMU receptor, which comprises separately contacting cells producing a second messenger response and expressing on their cell surface the mammalian NMU receptor, wherein such cells do not normally express the mammalian NMU receptor, with both the chemical compound and a second chemical compound known to activate the mammalian NMU receptor, and with only the second chemical compound, under conditions suitable for activation of the mammalian NMU receptor, and measuring the second messenger response in the presence of only the second chemical compound and in the presence of both the second chemical compound and the chemical compound, a smaller change, e.g. increase, in the second messenger response in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound indicating that the chemical compound inhibits activation of the mammalian NMU receptor.

In an embodiment of the present invention, the second chemical compound is an NMU peptide. Examples of NMU peptides include, but are not limited to, human NMU-25, human NMU-8, porcine NMU-8, porcine NMU-25, rat NMU-25 and any peptide comprising the carboxyl terminal seven amino acid residues of human NMU-8.

009509146.0630000  
5 In one embodiment, the second messenger response comprises chloride channel activation and the change in second messenger response is a smaller increase in the level of chloride current in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound. In another embodiment, the second messenger response comprises change in intracellular calcium levels and the change in second messenger response is a smaller increase in the measure of intracellular calcium in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound. In another embodiment, the second messenger response comprises release of inositol phosphate and the change in second messenger response is a smaller increase in the level of inositol phosphate in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound.

20 In one embodiment, the second messenger response comprises activation of MAP kinase and the change in second messenger response is a smaller increase in the level of MAP kinase activation in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound. In another embodiment, the second messenger response comprises change in cAMP levels and the change in second messenger response is a smaller change in the level of cAMP in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound. In another embodiment, the second messenger response comprises release of arachidonic acid and the change in second messenger response is an increase in the level of arachidonic acid levels in the presence of both the chemical compound and the second chemical compound than in the

presence of only the second chemical compound. In a further embodiment, the second messenger response comprises GTP $\gamma$ S ligand binding and the change in second messenger is a smaller increase in GTP $\gamma$ S ligand binding in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound.

In one embodiment, the mammalian NMU receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In a further embodiment, the mammalian NMU receptor is a human SNORF72 receptor or a rat SNORF72 receptor. In another embodiment, the cell is an insect cell. In another embodiment, the cell is a mammalian cell. In another embodiment, the mammalian cell is nonneuronal in origin. In another embodiment, the nonneuronal cell is a COS-7 cell, CHO cell, 293 human embryonic kidney cell, NIH-3T3 cell or LM(tk-) cell. In another embodiment, the compound is not previously known to bind to a mammalian NMU receptor.

Further, this invention provides a compound determined by a process according to this invention and a composition, for example, a pharmaceutical composition, which comprises an amount of a mammalian SNORF62, a mammalian SNORF72 or a mammalian NMU receptor agonist determined to be such by a process according to this invention effective to increase activity of a mammalian SNORF62, a mammalian SNORF72 or a mammalian NMU receptor and a carrier, for example, a pharmaceutically acceptable carrier. In one embodiment, the mammalian SNORF62, mammalian SNORF72 or mammalian NMU receptor agonist is not previously known.

This invention also provides a composition, for example, a pharmaceutical composition, which comprises an amount of a mammalian SNORF62, a mammalian SNORF72 or a mammalian NMU



receptor antagonist determined to be such by a process according to this invention, effective to reduce activity of the mammalian SNORF62, the mammalian SNORF72 or the mammalian NMU receptor and a carrier, for example a pharmaceutically acceptable carrier. In one embodiment, the mammalian SNORF62, mammalian SNORF72 or mammalian NMU receptor antagonist is not previously known.

This invention yet further provides a method of screening a plurality of chemical compounds not known to activate a mammalian SNORF62 receptor to identify a compound which activates the mammalian SNORF62 receptor which comprises: (a) contacting cells transfected with and expressing the mammalian SNORF62 receptor with the plurality of compounds not known to activate the mammalian SNORF62 receptor, under conditions permitting activation of the mammalian SNORF62 receptor; (b) determining whether the activity of the mammalian SNORF62 receptor is increased in the presence of one or more of the compounds; and if so (c) separately determining whether the activation of the mammalian SNORF62 receptor is increased by any compound included in the plurality of compounds, so as to thereby identify each compound which activates the mammalian SNORF62 receptor.

This invention yet further provides a method of screening a plurality of chemical compounds not known to activate a mammalian SNORF72 receptor to identify a compound which activates the mammalian SNORF72 receptor which comprises: (a) contacting cells transfected with and expressing the mammalian SNORF72 receptor with the plurality of compounds not known to activate the mammalian SNORF72 receptor, under conditions permitting activation of the mammalian SNORF72 receptor; (b) determining whether the activity of the mammalian SNORF72 receptor is increased in the presence of one or more of the compounds; and if so (c) separately

determining whether the activation of the mammalian SNORF72 receptor is increased by any compound included in the plurality of compounds, so as to thereby identify each compound which activates the mammalian SNORF72 receptor.

In an embodiment, the mammalian SNORF62 receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In a further embodiment, the mammalian SNORF72 receptor is a human SNORF72 receptor or a rat SNORF72 receptor.

This invention provides a method of screening a plurality of chemical compounds not known to inhibit the activation of a mammalian NMU receptor to identify a compound which inhibits the activation of the mammalian NMU receptor, which comprises: (a) contacting cells transfected with and expressing the mammalian NMU receptor with the plurality of compounds in the presence of a known mammalian NMU receptor agonist, under conditions permitting activation of the mammalian NMU receptor; (b) determining whether the extent or amount of activation of the mammalian NMU receptor is reduced in the presence of one or more of the compounds, relative to the extent or amount of activation of the mammalian NMU receptor in the absence of such one or more compounds; and if so (c) separately determining whether each such compound inhibits activation of the mammalian NMU receptor for each compound included in the plurality of compounds, so as to thereby identify any compound included in such plurality of compounds which inhibits the activation of the mammalian NMU receptor.

In one embodiment, the mammalian NMU receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In a further embodiment, the mammalian NMU receptor is a human SNORF72 receptor or a rat SNORF72

receptor. In another embodiment, wherein the cell is a mammalian cell. In another embodiment, the mammalian cell is non-neuronal in origin. In another embodiment, the non-neuronal cell is a COS-7 cell, a 293 human embryonic kidney cell, a LM(tk-) cell or an NIH-3T3 cell.

This invention also provides a composition, for example, a pharmaceutical composition, comprising a compound identified by a method according to this invention in an amount effective to increase mammalian NMU receptor activity and a carrier, for example, a pharmaceutically acceptable carrier.

This invention also provides a composition, for example, a pharmaceutical composition, comprising a compound identified by a method according to this invention in an amount effective to increase mammalian SNORF62 receptor activity and a carrier, for example, a pharmaceutically acceptable carrier.

This invention also provides a composition, for example, a pharmaceutical composition, comprising a compound identified by a method according to this invention in an amount effective to increase mammalian SNORF72 receptor activity and a carrier, for example, a pharmaceutically acceptable carrier.

This invention still further provides a composition, for example, a pharmaceutical composition, comprising a compound identified by a method according to this invention in an amount effective to decrease mammalian NMU receptor activity and a carrier, for example, a pharmaceutically acceptable carrier.

This invention still further provides a composition, for example, a pharmaceutical composition, comprising a compound

identified by a method according to this invention in an amount effective to decrease mammalian SNORF62 receptor activity and a carrier, for example, a pharmaceutically acceptable carrier.

This invention still further provides a composition, for example, a pharmaceutical composition, comprising a compound identified by a method according to this invention in an amount effective to decrease mammalian SNORF72 receptor activity and a carrier, for example, a pharmaceutically acceptable carrier.

Furthermore, this invention provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF62 receptor which comprises administering to the subject a compound which is a mammalian SNORF62 receptor agonist in an amount effective to treat the abnormality.

Furthermore, this invention provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF72 receptor which comprises administering to the subject a compound which is a mammalian SNORF72 receptor agonist in an amount effective to treat the abnormality.

In one embodiment, the abnormality is a regulation of a steroid hormone disorder, an epinephrine release disorder, a gastrointestinal disorder, a cardiovascular disorder, an electrolyte balance disorder, hypertension, diabetes, a respiratory disorder, asthma, a reproductive function disorder, an immune disorder, an endocrine disorder, a musculoskeletal disorder, a neuroendocrine disorder, a cognitive disorder, a memory disorder, somatosensory and neurotransmission disorders, metabolic disorders, a motor

coordination disorder, a sensory integration disorder, a motor integration disorder, a dopaminergic function disorder, an appetite disorder, such as anorexia or obesity, a sensory transmission disorder, drug addiction, an olfaction disorder, an autonomic nervous system disorder, pain, neuropsychiatric disorders, affective disorder, migraine, circadian disorders, visual disorders, urinary disorders, blood coagulation-related disorders, developmental disorders, or ischemia-reperfusion injury-related diseases.

In a further embodiment, the abnormality is Addison's disease, Cushing's disease or a stress-related disorder. In yet another embodiment, the compounds and/or compositions of the present invention may be used to prevent miscarriage, induce labor or treat dysmenorrhea.

This invention additionally provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF62 receptor which comprises administering to the subject a compound which is a mammalian SNORF62 receptor antagonist in an amount effective to treat the abnormality.

This invention additionally provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF72 receptor which comprises administering to the subject a compound which is a mammalian SNORF72 receptor antagonist in an amount effective to treat the abnormality.

In one embodiment, the abnormality is a regulation of a steroid hormone disorder, an epinephrine release disorder, a gastrointestinal disorder, a cardiovascular disorder, an electrolyte balance disorder, hypertension, diabetes, a

respiratory disorder, asthma, a reproductive function disorder, an immune disorder, an endocrine disorder, a musculoskeletal disorder, a neuroendocrine disorder, a cognitive disorder, a memory disorder, somatosensory and neurotransmission disorders, metabolic disorders, a motor coordination disorder, a sensory integration disorder, a motor integration disorder, a dopaminergic function disorder, an appetite disorder, such as anorexia or obesity, a sensory transmission disorder, drug addiction, an olfaction disorder, an autonomic nervous system disorder, pain, neuropsychiatric disorders, affective disorder, migraine, circadian disorders, visual disorders, urinary disorders, blood coagulation-related disorders, developmental disorders, or ischemia-reperfusion injury-related diseases.

In one embodiment, the mammalian NMU receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In another embodiment, the mammalian NMU receptor is a human SNORF72 receptor or a rat SNORF72 receptor.

This invention also provides a process for making a composition of matter which specifically binds to a mammalian NMU receptor which comprises identifying a chemical compound using a process in accordance with this invention and then synthesizing the chemical compound or a novel structural and functional analog or homolog thereof.

This invention also provides a process for making a composition of matter which specifically binds to a mammalian SNORF62 receptor which comprises identifying a chemical compound using a process in accordance with this invention and then synthesizing the chemical compound or a novel structural and functional analog or homolog thereof.

This invention also provides a process for making a composition of matter which specifically binds to a mammalian SNORF72 receptor which comprises identifying a chemical compound using a process in accordance with this invention and then synthesizing the chemical compound or a novel structural and functional analog or homolog thereof.

This invention further provides a process for preparing a composition, for example a pharmaceutical composition which comprises admixing a carrier, for example, a pharmaceutically acceptable carrier, and a pharmaceutically effective amount of a chemical compound identified by a process in accordance with this invention or a novel structural and functional analog or homolog thereof.

In one embodiment, the mammalian NMU receptor is a mammalian SNORF62 receptor. In another embodiment, the mammalian NMU receptor is a mammalian SNORF72 receptor. In a further embodiment, the mammalian SNORF62 receptor is a human SNORF62 receptor, a rat SNORF62a receptor or a rat SNORF62b receptor. In a further embodiment, the mammalian SNORF72 receptor is a human SNORF72 receptor or a rat SNORF72 receptor.

Thus, once the gene for a targeted receptor subtype is cloned, it is placed into a recipient cell which then expresses the targeted receptor subtype on its surface. This cell, which expresses a single population of the targeted human receptor subtype, is then propagated resulting in the establishment of a cell line. This cell line, which constitutes a drug discovery system, is used in two different types of assays: binding assays and functional assays. In binding assays, the affinity of a compound for both the receptor subtype that is the target of a particular drug discovery program and other receptor subtypes that

could be associated with side effects are measured. These measurements enable one to predict the potency of a compound, as well as the degree of selectivity that the compound has for the targeted receptor subtype over other receptor subtypes. The data obtained from binding assays also enable chemists to design compounds toward or away from one or more of the relevant subtypes, as appropriate, for optimal therapeutic efficacy. In functional assays, the nature of the response of the receptor subtype to the compound is determined. Data from the functional assays show whether the compound is acting to inhibit or enhance the activity of the receptor subtype, thus enabling pharmacologists to evaluate compounds rapidly at their ultimate human receptor subtypes targets permitting chemists to rationally design drugs that will be more effective and have fewer or substantially less severe side effects than existing drugs.

Approaches to designing and synthesizing receptor subtype-selective compounds are well known and include traditional medicinal chemistry and the newer technology of combinatorial chemistry, both of which are supported by computer-assisted molecular modeling. With such approaches, chemists and pharmacologists use their knowledge of the structures of the targeted receptor subtype and compounds determined to bind and/or activate or inhibit activation of the receptor subtype to design and synthesize structures that will have activity at these receptor subtypes.

Combinatorial chemistry involves automated synthesis of a variety of novel compounds by assembling them using different combinations of chemical building blocks. The use of combinatorial chemistry greatly accelerates the process of generating compounds. The resulting arrays of compounds are called libraries and are used to screen for compounds



("lead compounds") that demonstrate a sufficient level of activity at receptors of interest. Using combinatorial chemistry it is possible to synthesize "focused" libraries of compounds anticipated to be highly biased toward the receptor target of interest.

Once lead compounds are identified, whether through the use of combinatorial chemistry or traditional medicinal chemistry or otherwise, a variety of homologs and analogs are prepared to facilitate an understanding of the relationship between chemical structure and biological or functional activity. These studies define structure activity relationships which are then used to design drugs with improved potency, selectivity and pharmacokinetic properties. Combinatorial chemistry is also used to rapidly generate a variety of structures for lead optimization. Traditional medicinal chemistry, which involves the synthesis of compounds one at a time, is also used for further refinement and to generate compounds not accessible by automated techniques. Once such drugs are defined the production is scaled up using standard chemical manufacturing methodologies utilized throughout the pharmaceutical and chemistry industry.

This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

## EXPERIMENTAL DETAILS

### Materials and Methods

#### Isolation of a full-length human SNORF62 receptor

The SwissPlus database was searched for G protein-coupled receptor sequences (GPCRs) using a select set of known GPCRs and the Wisconsin Package (GCG, Genetics Computer Group, Madison, WI). One sequence, O43664, was found which was most similar to the neurotensin receptor 1 (31% identity), as well as the recently identified motilin receptor, GPR38 (33% identity). O43664 was then chosen to be cloned for use in ligand-identification screens.

Before cloning the full-length receptor, the 5' and 3' ends of the coding sequence were verified by 5'/3' Rapid Amplification of cDNA Ends (RACE), using human hypothalamic Marathon-Ready cDNA (Clontech, Palo Alto, CA), the Marathon adaptor primers AP1 and AP2 (Clontech), and the primers JAB620, JAB621, JAB623, JAB624 and JAB622 set forth below.

A band of approximately 550 bp from the 5' RACE reaction was isolated from an agarose gel using the Qiaquick gel extraction kit (QIAGEN, Valencia, CA) and sequenced with JAB622. The sequence of this band indicated that the actual coding sequence of this receptor is longer than the sequence represented in the public database by 69 bp, coding for an additional potential initiating methionine 23 amino acids upstream from the methionine indicated by O43664. In addition, this 5' RACE sequence included some 5' untranslated sequence and an in-frame stop codon upstream from the new methionine. Sequencing of 3' RACE products revealed a sequence identical to the 3' end of O43664, in addition to some 3' untranslated sequence. The new (longer) coding sequence was named SNORF62.

From the new 5' untranslated sequence upstream from the new methionine and the 3' untranslated sequence, new primers were designed to amplify the entire SNORF62 sequence from human stomach cDNA using the Expand Long PCR system (Roche Biochemicals, Indianapolis, IN). The primers JAB648 and, JAB627 were designed to incorporate restriction sites for subcloning into the expression vector pEXJ.T3T7. The resulting PCR product of approximately 1330 bp was digested with HindIII and BamHI and subcloned into the HindIII/BamHI site of pEXJ.T3T7. This construct of SNORF62 with the additional methionine, subcloned into pEXJ.T3T7 was named pEXJ.T3T7-hSNORF62-f.

Primers and probes used in the identification of SNORF62:

JAB620 = 5'-CCACGAAGATCAGCAGGTATGTGG-3' (SEQ ID NO: 9)  
JAB621 = 5'-GGCATGAACAGCTCTGTCTGCTGG-3' (SEQ ID NO: 10)  
JAB623 = 5'-CCAGCCGCTTCCGAGAGACCTTCC-3' (SEQ ID NO: 11)  
JAB624 = 5'-GCCTGCTGCCATCGCCTCAGACCC-3' (SEQ ID NO: 12)  
JAB622 = 5'-GCCCCAGGTACTTGAGTCTCAGTG-3' (SEQ ID NO: 13)  
JAB648 = 5'-ATCTATAAGCTTCGGAGGGTGAAGCCGGGGTCTC-3' (SEQ ID NO: 14)  
JAB627 = 5'-ATCTATGGATCCTCAGGATGGATCGGTCTCTTGCTG-3' (SEQ ID NO: 15)

**Isolation of the rat SNORF62a and rat SNORF62b receptors**

To obtain a fragment of the rat homologue of SNORF62, 100 ng of rat genomic DNA (Clontech, Palo Alto, CA) and 1 µl of rat testes QUICK clone cDNA (Clontech) were amplified with forward PCR primers corresponding to TM1 (BB1611) or the 2nd intracellular loop (BB1614) and a reverse primer corresponding to TM6 (BB1612) of the mouse SNORF62 (GenEMBL Database Accession Number AF044602). PCR was performed with the Expand Long Template PCR System (Roche Molecular Biochemicals, Indianapolis, IN) under the following conditions: 30 seconds at 94°C, 45 seconds at 49°C to 67.7°C,

2 minutes at 68°C for 40 cycles; with a pre- and post-incubation of 5 minutes at 94°C and 7 minutes at 68°C, respectively. Bands of 430 and 700 bp from 6 independent reactions were isolated from a TAE gel, purified using the GENECLAN SPIN Kit (BIO101, Carlsbad, CA) and sequenced using the ABI BigDye cycle sequencing protocol and ABI 377 sequencers (ABI, Foster City, CA). Sequences were analyzed using the Wisconsin Package (GCG, Genetics Computer Group, Madison, WI). A consensus sequence was determined for these 6 products.

#### 5' and 3' RACE

The full length rat SNORF62 sequence was determined utilizing the Clontech Marathon cDNA Amplification kit (Clontech, Palo Alto, CA) for 5'/3' Rapid Amplification of cDNA ends (RACE). Nested PCR were performed according to the Marathon cDNA Amplification protocol using Marathon-Ready rat spleen cDNA (Clontech). For 5' RACE, the initial PCR was performed with the supplier's Adaptor Primer 1 and BB1631, a reverse primer from TM3 of the consensus sequence described above. Two µl of the initial PCR reaction was re-amplified using the Adaptor Primer 2 and BB1630, a reverse primer from the 1st extracellular loop. PCR was performed with Advantage KlenTaq Polymerase (Clontech, Palo Alto, CA) under the following conditions: 5 minutes at 94°C; 5 cycles of 94°C for 30 seconds and 72°C for 3 minutes; 5 cycles of 94°C for 30 seconds and 70°C for 3 minutes; 25 cycles (initial PCR) or 18 cycles (nested PCR) of 94°C for 30 seconds and 68°C for 3 minutes; 68°C hold for 7 minutes, and 4°C hold until the products were ready for analysis. 400 and 800 bp fragments were isolated from a 1% agarose TAE gel using the GENECLAN SPIN Kit and sequenced as above.

A second reaction was performed for 5' RACE using Marathon-Ready rat spleen and testes cDNA (Clontech). The initial

PCR was performed with the supplier's Adaptor Primer 1 and BB1650, a reverse primer from TM1 of the RACE fragment described above. Two  $\mu$ ls of the initial PCR reaction was re-amplified using the Adaptor Primer 2 and BB1649, a reverse primer from the amino terminus. PCR was performed as described above. 300 and 700 bp fragments were isolated from a 1% agarose TAE gel using the GENECLEAN SPIN Kit and sequenced as above.

For 3' RACE, initial PCR was performed using Marathon-Ready rat spleen with the supplier's Adapter Primer 1 and BB1632, a forward primer from TM5 of the consensus sequence described above. Two  $\mu$ ls of this initial PCR reaction was re-amplified using Adaptor Primer 2 and BB1633, a forward primer from the 3rd intracellular loop. PCR was performed with Advantage KlenTaq Polymerase (Clontech, Palo Alto, CA) under the following conditions: 5 minutes at 94°C; 5 cycles of 94°C for 30 seconds and 72°C for 3 minutes; 5 cycles of 94°C for 30 seconds and 70°C for 3 minutes; 25 cycles (initial PCR) or 18 cycles (nested PCR) of 94°C for 30 seconds and 68°C for 3 minutes; 68°C hold for 7 minutes, and 4°C hold until the products were ready for analysis. A 1000 bp fragment was isolated from a 1% agarose TAE gel using the GENECLEAN SPIN Kit and sequenced as above.

Primers and probes used in the identification of rat SNORF62a and rat SNORF62b:

BB1611 = 5'-TAC CTG CTG ATC TTC GTG GTG GG- 3' (SEQ ID NO: 30)

BB1612 = 5'-CAG TGC AAA CAG CAT CTT GGT CAC- 3' (SEQ ID NO: 31)

BB1614 = 5'-TAT GTG GCC GTG GTG CGC CCA CTC C- 3' (SEQ ID NO: 32)

BB1630 = 5'-CCA CCT GCA CCC AGC TGG AAT GGG- 3' (SEQ ID

NO: 33)

BB1631 = 5'-ACT GAA GCC AGG CAG ACA GTC TCC- 3' (SEQ ID

NO: 34)

BB1632 = 5'-TGG TCA CCA TCA GTG TGC TGT ACC- 3' (SEQ ID

NO: 35)

BB1633 = 5'-TGC GGA GGG AGA GGA TGT TGC TCC- 3' (SEQ ID

NO: 36)

BB1649 = 5'-CCC AAG TAC TTC AGC CTC AGG TCC- 3' (SEQ ID

NO: 37)

BB1650 = 5'-GGT CAA CCC GTT GCC CAG AGT GCC- 3' (SEQ ID

NO: 38)

**Isolation of other species homologs of SNORF62 receptor cDNA**

A nucleic acid sequence encoding a SNORF62 receptor cDNA from other species may be isolated using standard molecular biology techniques and approaches such as those described below:

Approach #1: A genomic library (e.g., cosmid, phage, P1, BAC, YAC) generated from the species of interest may be screened with a <sup>32</sup>P-labeled oligonucleotide probe corresponding to a fragment of the human SNORF62 receptor whose sequence is shown in Figures 1A-1B to isolate a genomic clone. The full-length sequence may be obtained by sequencing this genomic clone. If one or more introns are present in the gene, the full-length intronless gene may be obtained from cDNA using standard molecular biology techniques. For example, a forward PCR primer designed in the 5'UT and a reverse PCR primer designed in the 3'UT may be used to amplify a full-length, intronless receptor from cDNA. Standard molecular biology techniques could be used to subclone this gene into a mammalian expression vector.

Approach #2: Standard molecular biology techniques may be used to screen commercial cDNA phage libraries of the

species of interest by hybridization under reduced stringency with a  $^{32}\text{P}$ -labeled oligonucleotide probe corresponding to a fragment of the sequences shown in Figures 1A-1B. One may isolate a full-length SNORF62 receptor by obtaining a plaque purified clone from the lambda libraries and then subjecting the clone to direct DNA sequencing. Alternatively, standard molecular biology techniques could be used to screen cDNA plasmid libraries by PCR amplification of library pools using primers designed against a partial species homolog sequence. A full-length clone may be isolated by Southern hybridization of colony lifts of positive pools with a  $^{32}\text{P}$ -oligonucleotide probe.

Approach #3: 3' and 5' RACE may be utilized to generate PCR products from cDNA derived from the species of interest expressing SNORF62 which contain the additional sequence of SNORF62. These RACE PCR products may then be sequenced to determine the additional sequence. This new sequence is then used to design a forward PCR primer in the 5'UT and a reverse primer in the 3'UT. These primers are then used to amplify a full-length SNORF62 clone from cDNA.

Examples of other species include, but are not limited to, mouse, dog, monkey, hamster and guinea pig.

#### **Isolation of a full-length human SNORF72 receptor**

A search of the public domain databases revealed an amino acid sequence that was 46% identical to the amino acid sequence of SNORF62. This sequence was given the name human SNORF72. Primers were designed against the 5'- and 3'- untranslated regions of SNORF72, with restriction sites incorporated for subcloning purposes. GSL42 is a forward primer in the 5' untranslated region with a NotI site incorporated into the 5' end of the primer, and GSL43 is a reverse primer in the 3' untranslated region with an XbaI

site incorporated into the 5' end of the primer. These primers were used to amplify the full-length sequence from human whole-brain cDNA using the Expand Long Template PCR system (Roche Biochemicals, Indianapolis, IN). Sequencing of several clones from independent PCR reactions indicated that the actual sequence of SNORF72 differed from the published sequence at five base positions, four of which changed the amino acid sequence of the receptor. This sequence-verified SNORF72 clone was subcloned into the NotI/XbaI site of the mammalian expression vector pEXJ.T3T7 and named pEXJ.T3T7-hSNORF72-f.

Primers and probes used in the identification of SNORF72:

GSL42 = 5'-ATCTATGCGGCCGCTTGAAACAGAGCCTCGTACC-3' (SEQ ID NO: 16)

GSL43 = 5'-ATCTATTCTAGAGTTGTAAGAGCCATTCTACCTC-3' (SEQ ID NO: 17)

#### **Isolation of a full-length rat SNORF72 receptor**

A pair of oligo primers, BB1606 and BB1607 (set forth below), were synthesized based on sequence of the human SNORF72 gene. A PCR reaction was performed using this primer pair on rat brain cDNA from Clontech. The PCR condition used was 95°C for 5 minutes for initial denaturation, 94°C for 30 seconds followed by 50°C for 30 seconds and 68°C for 90 seconds for total of 40 cycles, finished with extension at 68°C for 7 minutes. The PCR product was sequenced and another pair of oligo primers, BB1609 and BB1610 (set forth below), were synthesized based on the sequencing information obtained. An oligo(dT) primed cDNA library from rat hypothalamus was screened by PCR using BB1609 and BB1610 as primers, and three positive pools were found from the first 188 pools screened. The PCR condition used was: 94°C for 4 minutes for initial denaturation, 94°C for 30 seconds followed by 68°C for 90 seconds for total of 40 cycles,



finished with extension at 68°C for 7 minutes. After two rounds of sib-selection, colonies were plated and positive clones were screened by hybridization with a radiolabeled oligonucleotide probe KS2008. A positive clone containing a 3.5kb cDNA insert was isolated and found to contain the full coding region of rat SNORF72 by sequence analysis.

Primers and probes used in the identification of rat SNORF72:

BB1606 = 5'-TCTATGAGATGTGGCGCAACTACC-3' (SEQ ID NO: 39)  
BB1607 = 5'-AACACTAAGACCAAGACAAACAGC-3' (SEQ ID NO: 40)  
BB1609 = 5'-GTCACCACGGTTAGCGTAGAGCGC-3' (SEQ ID NO: 41)  
BB1610 = 5'-GAGGGTCTGTGAATATTCACAGCC-3' (SEQ ID NO: 42)  
KS2008 = 5'-CCCAACGGGTCCTCCGTACCTGGCTCAGCCACCTGCACAGTCACC-3' (SEQ ID NO: 43)

**Isolation of other species homologs of SNORF72 receptor cDNA**

A nucleic acid sequence encoding a SNORF72 receptor cDNA from other species may be isolated using standard molecular biology techniques and approaches such as those described below:

Approach #1: A genomic library (e.g., cosmid, phage, P1, BAC, YAC) generated from the species of interest may be screened with a <sup>32</sup>P-labeled oligonucleotide probe corresponding to a fragment of the human SNORF72 receptor whose sequence is shown in Figures 3A-3B to isolate a genomic clone. The full-length sequence may be obtained by sequencing this genomic clone. If one or more introns are present in the gene, the full-length intronless gene may be obtained from cDNA using standard molecular biology techniques. For example, a forward PCR primer designed in the 5'UT and a reverse PCR primer designed in the 3'UT may be used to amplify a full-length, intronless receptor from

cDNA. Standard molecular biology techniques could be used to subclone this gene into a mammalian expression vector.

5 Approach #2: Standard molecular biology techniques may be used to screen commercial cDNA phage libraries of the species of interest by hybridization under reduced stringency with a <sup>32</sup>P-labeled oligonucleotide probe corresponding to a fragment of the sequences shown in Figures 3A-3B. One may isolate a full-length SNORF72 receptor by obtaining a plaque purified clone from the lambda libraries and then subjecting the clone to direct DNA sequencing. Alternatively, standard molecular biology techniques could be used to screen cDNA plasmid libraries by PCR amplification of library pools using primers designed against a partial species homolog sequence. A full-length clone may be isolated by Southern hybridization of colony lifts of positive pools with a <sup>32</sup>P-oligonucleotide probe.

20 Approach #3: 3' and 5' RACE may be utilized to generate PCR products from cDNA derived from the species of interest expressing SNORF72 which contain the additional sequence of SNORF72. These RACE PCR products may then be sequenced to determine the additional sequence. This new sequence is then used to design a forward PCR primer in the 5'UT and a reverse primer in the 3'UT. These primers are then used to amplify a full-length SNORF72 clone from cDNA.

Examples of other species include, but are not limited to, mouse, dog, monkey, hamster and guinea pig.

#### Host cells

30 A broad variety of host cells can be used to study heterologously expressed proteins. These cells include but are not limited to mammalian cell lines such as; COS-7, CHO, LM(tk<sup>-</sup>), HEK293, etc.; insect cell lines such as; Sf9, Sf21,

35

Trichoplusia ni 5B-4, etc.; amphibian cells such as *Xenopus* oocytes; assorted yeast strains; assorted bacterial cell strains; and others. Culture conditions for each of these cell types is specific and is known to those familiar with the art.

COS-7 cells are grown on 150 mm plates in DMEM with supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin 100 µg/ml streptomycin) at 37°C, 5% CO<sub>2</sub>. Stock plates of COS-7 cells are trypsinized and split 1:6 every 3-4 days.

#### Transient expression

DNA encoding proteins to be studied can be transiently expressed in a variety of mammalian, insect, amphibian, yeast, bacterial and other cells lines by several transfection methods including but not limited to; calcium phosphate-mediated, DEAE-dextran mediated; liposomal-mediated, viral-mediated, electroporation-mediated, and microinjection delivery. Each of these methods may require optimization of assorted experimental parameters depending on the DNA, cell line, and the type of assay to be subsequently employed.

A typical protocol for the DEAE-dextran method as applied to COS-7 and HEK293 cells is described as follows. Cells to be used for transfection are split 24 hours prior to the transfection to provide flasks which are 70-80% confluent at the time of transfection. Briefly, 8 µg of receptor DNA plus 8 µg of any additional DNA needed (e.g. G<sub>α</sub> protein expression vector, reporter construct, antibiotic resistance marker, mock vector, etc.) are added to 9 ml of complete DMEM plus DEAE-dextran mixture (10 mg/ml in PBS). Cells plated into a T225 flask (sub-confluent) are washed once with PBS and the DNA mixture is added to each flask. The

cells are allowed to incubate for 30 minutes at 37°C, 5% CO<sub>2</sub>. Following the incubation, 36 ml of complete DMEM with 80 µM chloroquine is added to each flask and allowed to incubate an additional 3 hours. The medium is then aspirated and 24 ml of complete medium containing 10% DMSO for exactly 2 minutes and then aspirated. The cells are then washed 2 times with PBS and 30 ml of complete DMEM added to each flask. The cells are then allowed to incubate over night. The next day the cells are harvested by trypsinization and reseeded into 96 well plates.

#### Stable expression

Heterologous DNA can be stably incorporated into host cells, causing the cell to perpetually express a foreign protein. Methods for the delivery of the DNA into the cell are similar to those described above for transient expression but require the co-transfection of an ancillary gene to confer drug resistance on the targeted host cell. The ensuing drug resistance can be exploited to select and maintain cells that have taken up the DNA. An assortment of resistance genes are available including but not restricted to neomycin, kanamycin, and hygromycin. For purposes of studies concerning the receptor of this invention, stable expression of a heterologous receptor protein is typically carried out in, mammalian cells including but not necessarily restricted to, CHO, HEK293, LM(tk-), etc. In addition native cell lines that naturally carry and express the nucleic acid sequences for the receptor may be used without the need to engineer the receptor complement.

#### Functional assays

Cells expressing the receptor DNA of this invention may be used to screen for ligands to said receptor using functional assays. Once a ligand is identified the same assays may be used to identify agonists or antagonists of the receptor

that may be employed for a variety of therapeutic purposes. It is well known to those in the art that the over-expression of a G-protein coupled receptor can result in the constitutive activation of intracellular signaling pathways. In the same manner, over-expression of the receptors of the present invention in any cell line as described above, can result in the activation of the functional responses described below, and any of the assays herein described can be used to screen for agonist, partial agonist, inverse agonist and antagonist ligands of the SNORF62 and SNORF72 receptors.

A wide spectrum of assays can be employed to screen for the presence of receptor SNORF62 and SNORF72 ligands. These assays range from traditional measurements of total inositol phosphate accumulation, cAMP levels, intracellular calcium mobilization, and potassium currents, for example; to systems measuring these same second messengers but which have been modified or adapted to be of higher throughput, more generic and more sensitive; to cell based assays reporting more general cellular events resulting from receptor activation such as metabolic changes, differentiation, cell division/proliferation. Description of several such assays follow.

#### Cyclic AMP (cAMP) assay

The receptor-mediated stimulation or inhibition of cyclic AMP (cAMP) formation may be assayed in cells expressing the receptors. COS-7 cells are transiently transfected with the receptor gene using the DEAE-dextran method and plated in 96-well plates. 48 hours after transfection, cells are washed twice with Dulbecco's phosphate buffered saline (PBS) supplemented with 10 mM HEPES, 10 mM glucose and 5 mM theophylline and are incubated in the same buffer for 20 min at 37°C, in 5% CO<sub>2</sub>. Test compounds are added and cells are

incubated for an additional 10 min at 37°C. The medium is then aspirated and the reaction stopped by the addition of 100 mM HCl. The plates are stored at -20°C for 2-5 days. For cAMP measurement, plates are thawed and the cAMP content in each well is measured by cAMP Scintillation Proximity Assay (Amersham Pharmacia Biotech). Radioactivity is quantified using microbeta Trilux counter (Wallac).

#### Arachidonic acid release assay

Cells expressing the receptor are seeded into 96 well plates or other vessels and grown for 3 days in medium with supplements. <sup>3</sup>H-arachidonic acid (specific activity = 0.75 µCi/ml) is delivered as a 100 µL aliquot to each well and samples are incubated at 37° C, 5% CO<sub>2</sub> for 18 hours. The labeled cells are washed three times with medium. The wells are then filled with medium and the assay is initiated with the addition of test compounds or buffer in a total volume of 250 µL. Cells are incubated for 30 min at 37°C, 5% CO<sub>2</sub>. Supernatants are transferred to a microtiter plate and evaporated to dryness at 75°C in a vacuum oven. Samples are then dissolved and resuspended in 25 µL distilled water. Scintillant (300 µL) is added to each well and samples are counted for <sup>3</sup>H in a Trilux plate reader. Data are analyzed using nonlinear regression and statistical techniques available in the GraphPAD Prism package (San Diego, CA).

#### Intracellular calcium mobilization assays

The intracellular free calcium (Ca<sup>2+</sup>) concentration may be measured by microspectrofluorimetry using the fluorescent indicator dye Fura-2/AM (Bush et al., 1991). Cells expressing the receptor are seeded onto a 35mm culture dish containing a glass coverslip insert and allowed to adhere overnight. Cells are then washed with HBS and loaded with 100 µL of Fura-2/AM (10 µM) for 20 to 40 min. After washing with HBS to remove the Fura-2/AM solution, cells are

equilibrated in HBS for 10 to 20 min. Cells are then visualized under the 40X objective of a Leitz Fluovert FS microscope and fluorescence emission is determined at 510 nM with excitation wavelengths alternating between 340 nM and 380 nM. Raw fluorescence data are converted to  $\text{Ca}^{2+}$  concentrations using standard  $\text{Ca}^{2+}$  concentration curves and software analysis techniques.

In another method, the measurement of intracellular  $\text{Ca}^{2+}$  can also be performed on a 96-well (or higher) format and with alternative  $\text{Ca}^{2+}$ -sensitive indicators, preferred examples of these are: aequorin, Fluo-3, Fluo-4, Fluo-5, Calcium Green-1, Oregon Green, and 488 BAPTA. After activation of the receptors with agonist ligands the emission elicited by the change of intracellular  $\text{Ca}^{2+}$  concentration can be measured by a luminometer, or a fluorescence imager; a preferred example of this is the fluorescence imager plate reader (FLIPR™, Molecular Devices).

Cells expressing the receptor of interest are plated into clear, flat-bottom, black-walled 96-well plates (Costar) at a density of 80,000-150,000 cells per well and allowed to incubate for 48 hr at 5%  $\text{CO}_2$ , 37°C. The growth medium is aspirated and 100  $\mu\text{L}$  of loading medium containing Fluo-3 dye is added to each well. The loading medium contains: 20 mM HEPES (Sigma), 0.1% BSA (Sigma), dye/pluronic acid mixture (e.g. 1 mM Fluo-3/AM (Molecular Probes) and 10% pluronic acid (Molecular Probes) mixed immediately before use), and 2.5 mM probenecid (Sigma) (prepared fresh). The cells are allowed to incubate for about 1 hour at 5%  $\text{CO}_2$ , 37°C.

The compounds of interest are diluted in wash buffer (Hank's BSS (without phenol red), 20 mM HEPES, 2.5 mM probenecid) to a 4X final concentration and aliquoted into a clear v-bottom plate (Nunc). Following the dye incubation, the cells are

washed 4 times to remove excess dye using a Denley plate washer. 100  $\mu$ L final volume of wash buffer is then added to each cell well. Compounds are added to the cell plates and responses are measured using the FLIPR<sup>TM</sup> instrument. The data are then collected and analyzed using the FLIPR<sup>TM</sup> software and Graphpad Prism.

Antagonist ligands are identified by the inhibition of the signal elicited by agonist ligands.

#### GTP $\gamma$ S functional assay

Membranes from cells expressing the receptor are suspended in assay buffer (e.g., 50 mM Tris, 100 mM NaCl, 5 mM MgCl<sub>2</sub>, 10  $\mu$ M GDP, pH 7.4) with or without protease inhibitors (e.g., 0.1% bacitracin). Membranes are incubated on ice for 20 minutes, transferred to a 96-well Millipore microtiter GF/C filter plate and mixed with GTP $\gamma$ S<sup>35</sup> (e.g., 250,000 cpm/sample, specific activity ~1000 Ci/mmol) plus or minus unlabeled GTP $\gamma$ S (final concentration = 100  $\mu$ M). Final membrane protein concentration  $\approx$  90  $\mu$ g/ml. Samples are incubated in the presence or absence of test compounds for 30 min. at room temperature, then filtered on a Millipore vacuum manifold and washed three times with cold (4°C) assay buffer. Samples collected in the filter plate are treated with scintillant and counted for <sup>35</sup>S in a Trilux (Wallac) liquid scintillation counter. It is expected that optimal results are obtained when the receptor membrane preparation is derived from an appropriately engineered heterologous expression system, i.e., an expression system resulting in high levels of expression of the receptor and/or expressing G-proteins having high turnover rates (for the exchange of GDP for GTP). GTP $\gamma$ S assays are well-known to those skilled in the art, and it is contemplated that variations on the method described above, such as are described by Tian et al. (1994) or Lazareno and Birdsall (1993), may be used.



Microphysiometric assay

Because cellular metabolism is intricately involved in a broad range of cellular events (including receptor activation of multiple messenger pathways), the use of microphysiometric measurements of cell metabolism can in principle provide a generic assay of cellular activity arising from the activation of any orphan receptor regardless of the specifics of the receptor's signaling pathway.

General guidelines for transient receptor expression, cell preparation and microphysiometric recording are described elsewhere (Salon, J.A. and Owicki, J.A., 1996). Typically cells expressing receptors are harvested and seeded at  $3 \times 10^5$  cells per microphysiometer capsule in complete media 24 hours prior to an experiment. The media is replaced with serum free media 16 hours prior to recording to minimize non-specific metabolic stimulation by assorted and ill-defined serum factors. On the day of the experiment the cell capsules are transferred to the microphysiometer and allowed to equilibrate in recording media (low buffer RPMI 1640, no bicarbonate, no serum (Molecular Devices Corporation, Sunnyvale, CA) containing 0.1% fatty acid free BSA), during which a baseline measurement of basal metabolic activity is established.

A standard recording protocol specifies a 100  $\mu$ l/min flow rate, with a 2 min total pump cycle which includes a 30 sec flow interruption during which the acidification rate measurement is taken. Ligand challenges involve a 1 min 20 sec exposure to the sample just prior to the first post challenge rate measurement being taken, followed by two additional pump cycles for a total of 5 min 20 sec sample exposure. Typically, drugs in a primary screen are presented to the cells at 10  $\mu$ M final concentration.

Follow up experiments to examine dose-dependency of active compounds are then done by sequentially challenging the cells with a drug concentration range that exceeds the amount needed to generate responses ranging from threshold to maximal levels. Ligand samples are then washed out and the acidification rates reported are expressed as a percentage increase of the peak response over the baseline rate observed just prior to challenge.

#### MAP kinase assay

MAP kinase (mitogen activated kinase) may be monitored to evaluate receptor activation. MAP kinase is activated by multiple pathways in the cell. A primary mode of activation involves the ras/raf/MEK/MAP kinase pathway. Growth factor (tyrosine kinase) receptors feed into this pathway via SHC/Grb-2/SOS/ras. Gi coupled receptors are also known to activate ras and subsequently produce an activation of MAP kinase. Receptors that activate phospholipase C (such as Gq/G11-coupled) produce diacylglycerol (DAG) as a consequence of phosphatidyl inositol hydrolysis. DAG activates protein kinase C which in turn phosphorylates MAP kinase.

MAP kinase activation can be detected by several approaches. One approach is based on an evaluation of the phosphorylation state, either unphosphorylated (inactive) or phosphorylated (active). The phosphorylated protein has a slower mobility in SDS-PAGE and can therefore be compared with the unstimulated protein using Western blotting. Alternatively, antibodies specific for the phosphorylated protein are available (New England Biolabs) which can be used to detect an increase in the phosphorylated kinase. In either method, cells are stimulated with the test compound and then extracted with Laemmli buffer. The soluble fraction is applied to an SDS-PAGE gel and proteins are

transferred electrophoretically to nitrocellulose or Immobilon. Immunoreactive bands are detected by standard Western blotting technique. Visible or chemiluminescent signals are recorded on film and may be quantified by densitometry.

Another approach is based on evaluation of the MAP kinase activity via a phosphorylation assay. Cells are stimulated with the test compound and a soluble extract is prepared. The extract is incubated at 30°C for 10 min with gamma-<sup>32</sup>P-ATP, an ATP regenerating system, and a specific substrate for MAP kinase such as phosphorylated heat and acid stable protein regulated by insulin, or PHAS-I. The reaction is terminated by the addition of H<sub>3</sub>PO<sub>4</sub> and samples are transferred to ice. An aliquot is spotted onto Whatman P81 chromatography paper, which retains the phosphorylated protein. The chromatography paper is washed and counted for <sup>32</sup>P in a liquid scintillation counter. Alternatively, the cell extract is incubated with gamma-<sup>32</sup>P-ATP, an ATP regenerating system, and biotinylated myelin basic protein bound by streptavidin to a filter support. The myelin basic protein is a substrate for activated MAP kinase. The phosphorylation reaction is carried out for 10 min at 30°C. The extract can then be aspirated through the filter, which retains the phosphorylated myelin basic protein. The filter is washed and counted for <sup>32</sup>P by liquid scintillation counting.

#### Cell proliferation assay

Receptor activation of the orphan receptor may lead to a mitogenic or proliferative response which can be monitored via <sup>3</sup>H-thymidine uptake. When cultured cells are incubated with <sup>3</sup>H-thymidine, the thymidine translocates into the nuclei where it is phosphorylated to thymidine triphosphate. The nucleotide triphosphate is then incorporated into the

cellular DNA at a rate that is proportional to the rate of cell growth. Typically, cells are grown in culture for 1-3 days. Cells are forced into quiescence by the removal of serum for 24 hrs. A mitogenic agent is then added to the media. 24 hrs later, the cells are incubated with  $^3\text{H}$ -thymidine at specific activities ranging from 1 to 10  $\mu\text{Ci/ml}$  for 2-6 hrs. Harvesting procedures may involve trypsinization and trapping of cells by filtration over GF/C filters with or without a prior incubation in TCA to extract soluble thymidine. The filters are processed with scintillant and counted for  $^3\text{H}$  by liquid scintillation counting. Alternatively, adherent cells are fixed in MeOH or TCA, washed in water, and solubilized in 0.05% deoxycholate/0.1 N NaOH. The soluble extract is transferred to scintillation vials and counted for  $^3\text{H}$  by liquid scintillation counting.

Alternatively, cell proliferation can be assayed by measuring the expression of an endogenous or heterologous gene product, expressed by the cell line used to transfect the orphan receptor, which can be detected by methods such as, but not limited to, fluorescence intensity, enzymatic activity, immunoreactivity, DNA hybridization, polymerase chain reaction, etc.

#### Promiscuous second messenger assays

It is not possible to predict, a priori and based solely upon the GPCR sequence, which of the cell's many different signaling pathways any given receptor will naturally use.

It is possible, however, to coax receptors of different functional classes to signal through a pre-selected pathway through the use of promiscuous  $G_\alpha$  subunits. For example, by providing a cell based receptor assay system with an endogenously supplied promiscuous  $G_\alpha$  subunit such as  $G_{\alpha 15}$  or  $G_{\alpha 16}$  or a chimeric  $G_\alpha$  subunit such as  $G_{\alpha qz}$ , a GPCR, which might

normally prefer to couple through a specific signaling pathway (e.g.,  $G_s$ ,  $G_i$ ,  $G_q$ ,  $G_0$ , etc.), can be made to couple through the pathway defined by the promiscuous  $G_\alpha$  subunit and upon agonist activation produce the second messenger associated with that subunit's pathway. In the case of  $G_{\alpha 15}$ ,  $G_{\alpha 16}$  and/or  $G_{\alpha qz}$  this would involve activation of the  $G_q$  pathway and production of the second messenger  $IP_3$ . Through the use of similar strategies and tools, it is possible to bias receptor signaling through pathways producing other second messengers such as  $Ca^{++}$ , cAMP, and  $K^+$  currents, for example (Milligan and Rees, 1999).

It follows that the promiscuous interaction of the exogenously supplied  $G_\alpha$  subunit with the receptor alleviates the need to carry out a different assay for each possible signaling pathway and increases the chances of detecting a functional signal upon receptor activation.

#### Methods for recording currents in *Xenopus* oocytes

Oocytes are harvested from *Xenopus laevis* and injected with mRNA transcripts as previously described (Quick and Lester, 1994; Smith et al., 1997). The test receptor of this invention and  $G_\alpha$  subunit RNA transcripts are synthesized using the T7 polymerase ("Message Machine," Ambion) from linearized plasmids or PCR products containing the complete coding region of the genes. Oocytes are injected with 10 ng synthetic receptor RNA and incubated for 3-8 days at 17 degrees. Three to eight hours prior to recording, oocytes are injected with 500 pg promiscuous  $G_\alpha$  subunits mRNA in order to observe coupling to  $Ca^{++}$  activated  $Cl^-$  currents. Dual electrode voltage clamp (Axon Instruments Inc.) is performed using 3 M KCl-filled glass microelectrodes having resistances of 1-2 MOhm. Unless otherwise specified, oocytes are voltage clamped at a holding potential of -80 mV. During recordings, oocytes are bathed in continuously

flowing (1-3 ml/min) medium containing 96 mM NaCl, 2 mM KCl, 1.8 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub>, and 5 mM HEPES, pH 7.5 (ND96). Drugs are applied either by local perfusion from a 10 µl glass capillary tube fixed at a distance of 0.5 mm from the oocyte, or by switching from a series of gravity fed perfusion lines.

Other oocytes may be injected with a mixture of receptor mRNAs and synthetic mRNA encoding the genes for G-protein-activated inward rectifier channels (GIRK1 and GIRK4, U.S. Patent Nos. 5,734,021 and 5,728,535 or GIRK1 and GIRK2) or any other appropriate combination (see, e.g., Inanobe et al., 1999). Genes encoding G-protein inwardly rectifying K<sup>+</sup> (GIRK) channels 1,2 and 4 (GIRK1, GIRK2, and GIRK4) may be obtained by PCR using the published sequences (Kubo et al., 1993; Dascal et al., 1993; Krapivinsky et al., 1995 and 1995b) to derive appropriate 5' and 3' primers. Human heart or brain cDNA may be used as template together with appropriate primers.

Heterologous expression of GPCRs in *Xenopus* oocytes has been widely used to determine the identity of signaling pathways activated by agonist stimulation (Gundersen et al., 1983; Takahashi et al., 1987). Activation of the phospholipase C (PLC) pathway is assayed by applying a test compound in ND96 solution to oocytes previously injected with mRNA for the human SNORF62 and observing inward currents at a holding potential of approximately -80 mV. The appearance of currents that reverse at -25 mV and display other properties of the Ca<sup>++</sup>-activated Cl<sup>-</sup> channel is indicative of receptor-activation of PLC and release of IP<sub>3</sub> and intracellular Ca<sup>++</sup>. Such activation is exhibited by GPCRs that couple to G<sub>q</sub> or G<sub>11</sub>.

Measurement of inwardly rectifying K<sup>+</sup> (potassium) channel

(GIRK) activity may be monitored in oocytes that have been co-injected with mRNAs encoding the mammalian receptor plus GIRK subunits. GIRK gene products co-assemble to form a G-protein activated potassium channel known to be activated (i.e., stimulated) by a number of GPCRs that couple to  $G_i$  or  $G_o$  (Kubo et al., 1993; Dascal et al., 1993). Oocytes expressing the mammalian receptor plus the GIRK subunits are tested for test compound responsitivity by measuring  $K^+$  currents in elevated  $K^+$  solution containing 49 mM  $K^+$ .

In the present invention, oocytes were harvested from *Xenopus laevis* and injected with mRNA transcripts as previously described (Quick and Lester, 1994; Smith et al., 1997). SNORF62 mRNA transcript was synthesized using the T7 polymerase ("Message Machine", Ambion) from linearized plasmids or PCR products containing the complete coding region of the gene. Oocytes were injected with 1-50 ng synthetic receptor RNA and incubated for 3-8 days at 17°C. Currents were recorded under dual electrode voltage clamp (Axon Instruments Inc.) with 3 M KCl-filled glass microelectrodes having resistances of 1-2 Mohm. Unless otherwise specified, oocytes were voltage clamped at a holding potential of -80 mV. During recordings, oocytes were bathed in continuously flowing (1-3 mL/min) medium containing 96 mM NaCl, 2 mM KCl, 1.8 mM  $CaCl_2$ , 1 mM  $MgCl_2$ , and 5 mM HEPES, pH 7.5 (ND96). Drugs were applied either by local perfusion from a 10  $\mu$ L glass capillary tube fixed at a distance of 0.5 mm from the oocyte, or by switching from a series of gravity fed perfusion lines.

#### Inositol phosphate assay

Human SNORF62 receptor-mediated activation and human SNORF72 receptor-mediated activation of the inositol phosphate (IP) second messenger pathways were assessed by radiometric measurement of IP products.

For example, in a 96 well microplate format assay, COS-7 cells expressing the receptor of interest were plated at a density of 70,000 cells per well and allowed to incubate for 24 hours. The cells were then labeled with 0.5  $\mu$ Ci [ $^3$ H]myo-inositol overnight at 37°C, 5% CO<sub>2</sub>. Immediately before the assay, the medium was removed and replaced with 180  $\mu$ L of Phosphate-Buffered Saline (PBS) containing 10 mM LiCl. The plates were then incubated for 20 min at 37°C, 5% CO<sub>2</sub>.

Following the incubation, the cells were challenged with agonist (20  $\mu$ l/well; 10x concentration) for 30 min at 37°C. The challenge was terminated by the addition of 100  $\mu$ L of 5% v/v trichloroacetic acid, followed by incubation at 4°C for greater than 30 minutes. Total IPs were isolated from the lysate by ion exchange chromatography. Briefly, the lysed contents of the wells were transferred to a Multiscreen HV filter plate (Millipore) containing Dowex AG1-X8 (200-400 mesh, formate form). The filter plates were prepared adding 100  $\mu$ L of Dowex AG1-X8 suspension (50% v/v, water: resin) to each well. The filter plates were placed on a vacuum manifold to wash or elute the resin bed. Each well was first washed 2 times with 200  $\mu$ l of 5 mM myo-inositol. Total [ $^3$ H]inositol phosphates were eluted with 75  $\mu$ l of 1.2M ammonium formate/0.1M formic acid solution into 96-well plates. 200  $\mu$ L of scintillation cocktail was added to each well, and the radioactivity was determined by liquid scintillation counting.

#### Membrane preparations

Cell membranes expressing the receptor protein of this invention are useful for certain types of assays including but not restricted to ligand binding assays, GTP- $\gamma$ -S binding assays, and others. The specifics of preparing such cell membranes may in some cases be determined by the nature of the ensuing assay but typically involve harvesting whole cells and disrupting the cell pellet by sonication in ice



5

## 5

20

30

## 35

Cells expressing the receptors of this invention may be used to screen for ligands for said receptors, for example, by

[<sup>125</sup>I]rat NMU-23 and [<sup>125</sup>I]NMU-8 binding assays. The same assays may be used to identify agonists or antagonists of the receptor that may be employed for a variety of therapeutic purposes.

5 Radioligand binding assays were performed by diluting membranes prepared from cells expressing the receptor in 50 mM Tris buffer (pH = 7.4 at 0°C) containing 0.1% bovine serum albumin (Sigma), aprotinin (0.005 mg/ml, Boehringer Mannheim) and bestatin (0.1 mM, Sigma) as protease inhibitors. The final protein concentration in the assay was 12 - 40 µg/ml. Membranes were then incubated with either [<sup>125</sup>I]rat NMU-23 or [<sup>125</sup>I]NMU-8 (NEN, specific activity 2200 Ci/mmol) in the presence or absence of competing ligands on ice for 60 min in a total volume of 250 µl in 96 well microtiter plates. The bound ligand was separated from free by filtration through GF/B filters presoaked in 0.5% polyethyleneimine (PEI), using a Tomtec (Wallac) vacuum filtration device. After addition of Ready Safe (Beckman) scintillation fluid, bound radioactivity was quantitated using a Trilux (Wallac) scintillation counter (approximately 40% counting efficiency of bound counts). Data was fit to non-linear curves using GraphPad Prism.

25 In this manner, agonist or antagonist compounds that bind to the receptor may be identified as they inhibit the binding of the labeled ligand to the membrane protein of cells expressing the said receptor. Non-specific binding was defined as the amount of radioactivity remaining after incubation of membrane protein in the presence of 100 nM of the unlabeled peptide corresponding to the radioligand used. In equilibrium saturation binding assays membrane preparations or intact cells transfected with the receptor are incubated in the presence of increasing concentrations of the labeled compound to determine the binding affinity of

the labeled ligand. The binding affinities of unlabeled compounds may be determined in equilibrium competition binding assays, using a fixed concentration of labeled compound (0.05 - 0.1 nM for [<sup>125</sup>I]rat NMU-23) in the presence of varying concentrations of the displacing ligands.

Localization of mRNA coding for human SNORF62 and human SNORF72.

Quantitative PCR using a fluorogenic probe with real time detection

Quantitative PCR using fluorogenic probes used to characterize the distribution of SNORF62 and SNORF72 RNA. This assay utilizes two oligonucleotides for conventional PCR amplification and a third specific oligonucleotide probe that is labeled with a reporter at the 5' end and a quencher at the 3' end of the oligonucleotide. In the instant invention, FAM (6-carboxyfluorescein) was used as the reporter, and TAMRA (6-carboxy-4,7,2,7'-tetramethyl-rhodamine) was the quencher. As amplification progresses, the labeled oligonucleotide probe hybridizes to the gene sequence between the two oligonucleotides used for amplification. The nuclease activity of *Taq* thermostable DNA polymerase is utilized to cleave the labeled probe. This separates the quencher from the reporter and generates a fluorescent signal that is directly proportional to the amount of amplicon generated. This labeled probe confers a high degree of specificity. Non-specific amplification is not detected as the labeled probe does not hybridize and as a consequence is not cleaved. All experiments were conducted in a PE7700 Sequence Detection System (Perkin Elmer, Foster City CA).

Quantitative RT-PCR

Quantitative RT-PCR was used for the detection of SNORF62 and SNORF72 RNA. For use as a template in quantitative PCR

reactions, cDNA was synthesized by reverse transcription from total human RNA. Reverse transcription by SuperScriptII RNase H<sup>-</sup> (GibcoBRL/life Technologies) was primed using random hexamers. Parallel reactions included <sup>32</sup>P labeled dCTP to allow quantification of the cDNA.

Following reverse transcription, cDNA was phenol/chloroform extracted and precipitated. Incorporation of <sup>32</sup>P dCTP was assessed after precipitation with trichloroacetic acid and the amount of cDNA synthesized was calculated.

For PCR reactions primers with the following oligonucleotide sequences were used:

**Human SNORF62:**

Forward primer

snorf62h.txt-115F

5'-CAATGGCAGTGCGGCC-3' (SEQ ID NO: 18)

Reverse primer

snorf62h.txt-239R

5'-GGTATGTGGCACAGATGGGC-3' (SEQ ID NO: 19)

Fluorogenic oligonucleotide probe:

snorf62h.txt-138T

5' (6-FAM)- ACTTTGACCCTGAGGACTTGAACCTGACTG-(TAMRA) 3' (SEQ ID NO: 20)

**Human SNORF72:**

Forward primer:

snorf 72h.txt-179F

5'-CCTCGGCGCAGCCAC-3' (SEQ ID NO: 21)

Reverse primer

snorf 72h.txt-275R

5'-GAATCACCAGGCACACCAGG-3' (SEQ ID NO: 22)

Fluorogenic oligonucleotide probe:

snorf 72h.txt-203T

5' (6-FAM)-CCCGTGTCTGTGGTGTATGTGCCAAT-(TAMRA) 3' (SEQ ID NO:  
23)

Using these primer pairs, amplicon length is 124 bp for  
SNORF62, and 96 bp for SNORF72. Each PCR reaction contained  
3.0 ng cDNA. Oligonucleotide concentrations were: 500 nM of  
forward and reverse primers, and 200 nM of fluorogenic  
probe. PCR reactions were carried out in 50 µl volumes  
using TaqMan universal PCR master mix (PE Applied  
Biosystems). Buffer for RT-PCR reactions contained a fluor  
used as a passive reference (ROX: Perkin Elmer proprietary  
passive reference I). All reagents for PCR (except cDNA and  
oligonucleotide primers) were obtained from Perkin Elmer  
(Foster City, CA). Reactions were carried in a PE7700  
sequence detection system (PE Applied Biosystems) using the  
following thermal cycler profile: 50°C 2 min., 95°C 10 min.,  
followed by 40 cycles of: 95°C, 15 sec., 60°C 1 min.

Positive controls for PCR reactions consisted of  
amplification of the target sequence from a plasmid  
construct when available. Standard curves for quantitation  
of human SNORF62 and SNORF72 were constructed using genomic  
DNA. Negative controls consisted of mRNA blanks, as well as  
primer and mRNA blanks. To confirm that the mRNA was not  
contaminated with genomic DNA, PCR reactions were carried  
out without reverse transcription using Taq DNA polymerase.  
Integrity of RNA was assessed by amplification of RNA coding  
for cyclophilin or glyceraldehyde 3-phosphate dehydrogenase  
(GAPDH). Following reverse transcription and PCR  
amplification, data was analyzed using Perkin Elmer sequence  
detection software. The fluorescent signal from each well  
was normalized using an internal passive reference, and data  
was fitted a standard curve to obtain relative quantities of

SNORF62 and SNORF72 expression.

Chromosomal localization of human SNORF62 and SNORF72  
receptor genes

5 Chromosomal localization for human SNORF62 and SNORF72  
receptor genes was established using a panel of radiation  
hybrids prepared by the Stanford Human Genome Center (SHGC)  
and distributed by Research Genetics, Inc. The Stanford G3  
panel of 83 radiation hybrids was analyzed by PCR using the  
same primers, probes and thermal cycler profiles as used for  
localization. 20 ng of DNA was used in each PCR reaction.  
Data was submitted to the RH Server (SHGC) which linked the  
SNORF62 and SNORF72 gene sequences to specific markers.  
NCBI LocusLink and NCBI GeneMap '99 were used to further  
analyze the data.

RT-PCR

For the detection of RNA encoding rat SNORF72, RT-PCR was  
carried out on mRNA extracted from tissue. Reverse  
transcription and PCR reactions were carried out in a 50 µl  
volumes using rTth DNA polymerase (Perkin Elmer). Primers  
with the following sequences were used:

rat SNORF72:

Forward primer:

snorf72rseq.txt-392F

5'-GCCTGTGGGATGCTACTTCAAG-3' (SEQ ID NO: 44)

Reverse primer

snorf72rseq.txt-471R

5'-CGCTAACCGTGGTGACACTG-3' (SEQ ID NO: 45)

Fluorogenic oligonucleotide probe:

snorf72rseq.txt-422T

5' (6-FAM)-CTTCGAGACTGTGTGCTTTGCCTCCATTC-(TAMRA) 3' (SEQ ID NO:

46)

Using these primer pairs, amplicon length is 79 bp for rat SNORF72. Each RT-PCR reaction contained 100 ng total RNA. Oligonucleotide concentrations were: 500 nM of forward and reverse primers, and 200 nM of fluorogenic probe. Concentrations of reagents in each reaction were: 300  $\mu$ M each of dGTP; dATP; dCTP; 600  $\mu$ M UTP; 3.0mM Mn(OAc)<sub>2</sub> ; 50 mM Bicine; 115 mM potassium acetate, 8% glycerol, 5 units rTth DNA polymerase, and 0.5 units of uracil N-glycosylase. Buffer for RT-PCR reactions also contained a fluor used as a passive reference (ROX: Perkin Elmer proprietary passive reference I). All reagents for RT-PCR (except mRNA and oligonucleotide primers) were obtained from Perkin Elmer (Foster City, CA). Reactions were carried using the following thermal cycler profile: 50°C 2 min., 60°C 30 min., 95°C 5 min., followed by 40 cycles of: 94°C, 20 sec., 62°C 1 min.

Standard curves for quantitation of rat SNORF72 were constructed using genomic DNA. Negative controls consisted of mRNA blanks. To confirm that the mRNA was not contaminated with genomic DNA, PCR reactions were carried out without reverse transcription using Taq polymerase. Integrity of RNA was assessed by amplification of mRNA coding for cyclophilin or glyceraldehyde 3-phosphate dehydrogenase (GAPDH). Following reverse transcription and PCR amplification, data was analyzed using Perkin Elmer sequence detection software. The fluorescent signal from each well was normalized using an internal passive reference, and data was fitted to a standard curve to obtain relative quantities of SNORF72 RNA expression.

## RESULTS AND DISCUSSION

### Isolation of a full-length human SNORF62 receptor

5 A search of the SwissPlus database with a search set of known  
GPCRs yielded several orphan GPCR sequences. One sequence,  
043664, was found to be most similar to the neurotensin  
receptor 1 (31% identity), as well as the recently identified  
motilin receptor, GPR38 (33% identity). 043664 was then chosen  
to be cloned for use in ligand-identification screens. In the  
process of verifying the 5' and 3' ends of the coding sequence  
for 043664 by RACE, an additional methionine was found upstream  
from the initiating methionine of 043664, which was in-frame  
with the rest of the sequence. This new receptor sequence  
would be 69 bp longer, potentially coding for a protein 23  
amino acids longer than 043664. This new sequence was named  
SNORF62, and is represented in Figures 1A-1B and Figures 2A-2B.  
The SNORF62 cDNA codes for a protein of 426 amino acids  
(Figures 2A-2B). There are three potential N-linked  
glycosylation sites in the extracellular N-terminal domain at  
amino acid positions 7, 27, and 41. The C-terminal tail  
contains two potential casein kinase II phosphorylation sites  
at threonines 366 and 397, and one potential protein kinase C  
phosphorylation site at serine 360.

### Isolation of the rat SNORF62a and rat SNORF62b receptors

25 A fragment of the rat homologue of SNORF62 was amplified from  
rat genomic DNA and rat testes cDNA by low stringency PCR using  
oligonucleotide primers designed against the mouse SNORF62  
(GenEMBL Accession Number AF044602). This fragment contains  
30 667 nucleotides of rat SNORF62, from the end of TM1 to the  
beginning of TM6.

To obtain the full-length rat SNORF62, 5' RACE was performed  
on rat spleen and rat testes, and 3' RACE was performed on rat  
35 spleen. The 5' RACE reaction yielded 400 and 800 bp bands that



contained sequence information through the first transmembrane domain to the amino terminus, but had no putative in-frame initiating methionine-coding sequence. A second 5' RACE reaction yielded 300 bp band from rat testes cDNA that contained sequence information through the first transmembrane domain and a putative in-frame initiating methionine-coding sequence. Another band of 700 bp from rat spleen cDNA yielded a different sequence containing sequence information through the first transmembrane domain and a putative in-frame initiating methionine-coding sequence. The 3' RACE reaction yielded a 1000 bp band that contained sequence for an in-frame stop codon downstream from the region coding for the seventh transmembrane domain.

Two full-length receptor sequences were identified and named SNORF62a (from rat testes) and SNORF62b (from rat spleen). These sequences are identical except for the first two amino acids of SNORF62a and the first 28 amino acids of SNORF62b. The largest open reading frame is 1239 and 1317 nucleotides (Figures 17A-B and 19A-B), and predicts a protein of 413 and 439 amino acids (Figures 18A-B and 20A-B) for rat SNORF62a and rat SNORF62b, respectively. A comparison of the rat SNORF62a and rat SNORF62b sequences with the human SNORF62 sequence reveals 76.5% and 75% nucleotide identities and 71% and 69.5% amino acid identities, respectively. An amino acid alignment of the sequences of SNORF62 is shown in Figures 21A-21C. Hydrophobicity (Kyte-Doolittle) analysis of the amino acid sequence of the full length clones indicates the presence of seven hydrophobic regions, which is consistent with the seven transmembrane domains of a G protein coupled receptor (Figures 18A-B and 20A-B).

#### Isolation of a full-length human SNORF72 receptor

A receptor sequence that was 46% identical to SNORF62 was found in the public domain, and subsequently named SNORF72. After

cloning the full-length receptor from human whole-brain cDNA by PCR, the actual sequence of SNORF72 was found to be slightly different from the published clone. Five nucleotide differences were discovered between this new SNORF72 sequence and the corresponding published sequence, four of which changed the amino acid sequence of the receptor. The new clone in the pEXJ.T3T7 expression vector was named pEXJ.T3T7-hSNORF72-f. The nucleotide sequence of SNORF72 is shown in Figure 3A-3B, and the predicted amino acid sequence of the receptor encoded by SNORF72 is shown in Figure 4A-4B. A GAP comparison of the amino acid sequences (Wisconsin Package version 10.0, Genetics Computer Group, Madison, WI) of SNORF72 with SNORF62 indicates that there is a 47% amino acid identity between the two receptors (Figure 5), suggesting that they are likely to be members of the same receptor subfamily.

#### Isolation of a full-length rat SNORF72 receptor

Sequencing of the cDNA insert shows a long open reading frame containing the full 1185 base pair coding region, corresponding to a predicted protein of 395 amino acids. Hydrophobicity analysis reveals the seven predicted transmembrane domains typical of G protein-coupled receptors. Sequence comparison with the predicted human SNORF72 coding region reveals 81% identity at the nucleotide level and 79% identity at the amino acid level. The rat SNORF72 N-terminus has a 5 amino acid deletion compared with human SNORF72 and contains an additional downstream methionine not present in human SNORF72 (Figures 16A-16B). Conversely, the human SNORF72 N-terminus contains an additional upstream methionine (Figures 16A-16B) not found in rat SNORF72. The C-terminus of rat SNORF72 also differs from human SNORF72 in that it is shorter by 12 amino acids (Figures 16A-16B). Rat and human SNORF72 were also compared with human SNORF62 (Figures 16A-16B). Rat and human SNORF72 show 47-49% amino acid identity with human SNORF62, a typical level of homology for receptor subtypes activated by the same

ligand. This construct of rat SNORF72 was named pEXJ.BS-rSNORF72-f.

5 Increase in intracellular  $\text{Ca}^{2+}$  release

0 COS-7 cells were transiently transfected with SNORF62, SNORF72  
or vector DNA (mock) as described in Materials and Methods.  
Application of human NMU-25 resulted in concentration-dependent  
release of intracellular  $\text{Ca}^{2+}$  (as measured by FLIPR™) in COS-7  
cells transfected with SNORF62 or SNORF72 (Figures 6 and 7).  
In contrast, human NMU-25 had no significant effect on  
intracellular  $\text{Ca}^{2+}$  release in vector-transfected cells (Figure  
6). The EC50 values obtained for the stimulation of SNORF62  
and SNORF72 by NMU and related peptides are listed in Table 1  
and Table 1A.

5 SNORF72-transfected COS-7 cells were also stimulated by rat  
NMU-23, NMU-8 and human NMU-25 (Figure 8). All of the peptides  
that activated SNORF62 and SNORF72 produced similar maximum  
responses and were therefore full agonists (data not shown).

20 The high potency of NMU-induced stimulation of SNORF62 and  
SNORF72 provides support for classifying these receptors as NMU  
receptors. For comparison, in isolated rat uterus preparations  
the EC50 concentration for contraction by rat NMU-23 is 0.2 nM  
25 (Domin et al. 1989). The slightly lower potency of rat NMU-23  
observed in the SNORF62- and SNORF72- transfected COS-7 cells  
(average EC50 = 2.1 nM for SNORF62 and 1.7, 5.1 nM; n=2, for  
SNORF72) may be due to species differences in the peptides  
and/or receptors as well as the artificial cell hosts.

30 NMU-8 and rat NMU-23 produced robust increases in intracellular  
 $\text{Ca}^{2+}$  (as measured by FLIPR™) in COS-7 cells transiently  
transfected with rat SNORF72 (Figure 22). The responses were  
dose dependent reaching maximal stimulation at concentrations  
35



**TABLE 1A**

Potency of NMU and related peptides for stimulation of  $\text{Ca}^{2+}$  release and binding affinities were determined in SNORF62 and SNORF72-transfected COS-7 cells (n=2-7). "N.D." = not determined.

	SNORF62		SNORF72	
	Avg EC50 (nM)	Avg Ki (nM)	Avg EC50 (nM)	Avg Ki (nM)
human NMU-25	4.0 ± 1.3	2.0 ± 0.4	2.4 ± 0.6	3.0 ± 0
porcine NMU-25	5.2 ± 0.8	1.1 ± 0.2	3.0 ± 0.7	1.8 ± 0.6
rat NMU-23	2.1 ± 0.5	0.5 ± 0.2	5.0 ± 1.7	0.5 ± 0.6
porcine NMU-8	1.1 ± 0.4	3.0 ± 1.3	1.2 ± 0.3	1.2 ± 0.3
frog PP*	N.D.	> 10 $\mu\text{M}$	N.D.	> 10 $\mu\text{M}$
rat PP	N.D.	> 10 $\mu\text{M}$	N.D.	> 10 $\mu\text{M}$
VIP <sup>†</sup>	N.D.	> 10 $\mu\text{M}$	N.D.	> 10 $\mu\text{M}$

\* pancreatic polypeptide

+ vasoactive intestinal peptide

#### Radioligand binding

Receptor binding was performed on SNORF62-, SNORF72- and mock-transfected COS-7 membranes using [ $^{125}\text{I}$ ]rat NMU-23 and [ $^{125}\text{I}$ ]NMU-8 as radioligands as described in the Methods.

Binding of [ $^{125}\text{I}$ ]rat NMU-23 and [ $^{125}\text{I}$ ]NMU-8 to the SNORF62 and SNORF72 membranes was time dependent (reaching equilibrium by 30 min, data not shown) and saturable (Figures 9A-9B and 11A-11B). No saturable, specific binding sites for either radioligand were present in the mock-transfected COS-7 cell membranes (data not shown).

In membranes from SNORF62-transfected COS-7 cells [ $^{125}\text{I}$ ]rat NMU-23 and [ $^{125}\text{I}$ ]NMU-8 bound with high affinity ( $K_d$  = 0.61, 0.72 nM and  $K_d$  = 1.2, 2.8 nM, respectively; n=2). See Figures 9A and 9B, respectively. [ $^{125}\text{I}$ ]NMU-8 identified 3-fold fewer sites than did [ $^{125}\text{I}$ ]rat NMU-23 ( $B_{\text{max}}$  = 3.8, 3.4 pmol/mg protein and  $B_{\text{max}}$  = 16.5, 9.9 pmol/mg protein, respectively; n=2) even

though NMU-8 and rat NMU-23 are both full agonists in the functional assay (data not shown). This may be due to technical limitations in reaching high enough concentrations of [ $^{125}$ I]NMU-8 to fully saturate the binding sites, since it demonstrates somewhat lower affinity than [ $^{125}$ I]rat NMU-23 (See Table 1). Non-specific binding represented approximately 6% and 30% of total binding for [ $^{125}$ I]rat NMU-23 and [ $^{125}$ I]NMU-8, respectively.

In SNORF72-transfected COS-7 membranes, Kd values determined from saturation binding were 0.81, 0.96 nM (n=2) for [ $^{125}$ I]rat NMU-23 and Kd = 0.83, 0.82 (n=2) for [ $^{125}$ I]NMU-8. See Figures 11A and 11B, respectively. Non-specific binding represented approximately 15% and approximately 35% of total binding for [ $^{125}$ I]rat NMU-23 and [ $^{125}$ I]NMU-8, respectively. Both radioligands identified similar numbers of sites in SNORF72-transfected membranes (Bmax = 8.0, 8.2 pmol/mg protein for [ $^{125}$ I]rat NMU-23 and Bmax = 6.9, 5.5 pmol/mg protein for [ $^{125}$ I]NMU-8; n=2). Interestingly, although the maximum Ca<sup>2+</sup> signal generated by human NMU-25 was higher in SNORF72-transfected cells than in SNORF62-transfected cells (Figures 7 and 8 vs Figure 6), the levels of receptor expression were similar (based on Bmax values discussed above). This observation suggests more efficient coupling of SNORF72 to Ca<sup>2+</sup> releasing signal transduction mechanisms in the COS-7 cells.

Displacement of [ $^{125}$ I]rat NMU-23 binding allowed the estimation of binding affinity of a number of peptides (Figures 10, 12 and Table 2). The binding affinity of rat NMU-23 (average pKi = 9.4) in SNORF62 membranes was somewhat greater than NMU-8 (average pKi = 8.5), consistent with the binding affinities determined for the radioiodinated peptides. However, in SNORF72 membranes the apparent binding affinities of rat NMU-23 and NMU-8 were similar (average pKi = 9.0 and 8.9, respectively).

The high affinity binding of [<sup>125</sup>I]rat NMU-23 at SNORF62 and SNORF72 in COS-7 cell membranes is similar to the K<sub>d</sub> determined for this radioligand in isolated rat uterus (K<sub>d</sub> = 0.35 nM, Nandha et al. 1993). This affinity corresponds to the EC<sub>50</sub> of contractile activity in this tissue, 0.2nM, similar to the involvement of the binding site in NMU-induced uterine contraction. The binding site in this tissue demonstrated lower affinity for NMU-8 than for rat NMU-23 with average IC<sub>50</sub> values of 60 nM and 1 nM, respectively.

Although rat PP, frog PP and vasoactive intestinal peptide (VIP) share minor homology with NMU (see Background), these peptides did not displace binding of [<sup>125</sup>I]rat NMU-23 bound in either SNORF62 or SNORF72 membranes (Table 2) or activate the receptors in transfected COS-7 cells (data not shown).

**TABLE 2**

Binding pK<sub>i</sub> values determined from displacement of [<sup>125</sup>I]rat NMU-23 (0.05 - 0.1 nM) binding in membranes prepared from SNORF62- or SNORF72- transfected COS-7 cells. n=2, N.D. = not determined.

Compound	SNORF62		SNORF72	
	Avg pK <sub>i</sub>	STDev pK <sub>i</sub>	Avg pK <sub>i</sub>	STDev pK <sub>i</sub>
human NMU-25	8.7	0.1	8.5	0.1
porcine NMU-25	9.0	0.1	8.7	0.1
rat NMU-23	9.4	0.2	9.0	0.6
NMU-8	8.5	0.2	8.9	0.1
frog PP	<5	<5	<5	<5
rat PP	<5	<5	<5	<5
VIP	<5	<5	<5	<5

Inositol phosphate (IP) release

Exposure of SNORF62-transfected COS-7 cells (but not mock-transfected cells) to human NMU-25 caused the dose-dependent release of IP second messengers (approximately 2-fold above basal) with an EC50 of  $0.25 \pm 0.09$  nM (Figure 23). Rat NMU-23 and porcine NMU-8 were also full agonists in this assay with an EC50 =  $0.23 \pm 0.10$  nM and  $0.23 \pm 0.06$  nM, respectively (n=3). The EC50 values measured for IP release are lower than the EC50 values measured for increases in intracellular  $\text{Ca}^{2+}$  (see Table 1 and Table 1A). This may be due to differences in experimental conditions between the two types of assays including the non-equilibrium nature of  $\text{Ca}^{2+}$  measurements.

In addition, the  $\text{Ca}^{2+}$  release response to human NMU-25 was present in SNORF62-transfected cells following pretreatment with pertussis toxin (100 ng/ml for 18-20 hours, n=2) indicating that the  $\text{Ca}^{2+}$  signal is not predominantly generated by G-proteins of the Gi/Go family (data not shown). Taken together these results indicate that SNORF62 couples to phospholipase C stimulation via a Gq-type G-protein in COS-7 cells.

Activation of calcium-activated  $\text{Cl}^-$  currents in human SNORF62-expressing *Xenopus* oocytes

In *Xenopus laevis* oocytes injected with SNORF62 mRNA human NMU-25 elicited oscillatory  $\text{Cl}^-$  currents through G protein-coupled stimulation of the phosphoinositide/ $\text{Ca}^{2+}$  second messenger system, which in turn leads to the activation of a  $\text{Ca}^{2+}$ -dependent  $\text{Cl}^-$  current. As shown in Figure 13, control oocytes, lacking injection of foreign mRNA, typically showed no response to human NMU-25 (n=5). However, in oocytes injected with SNORF62 mRNA, the current amplitude averaged  $1065 \pm 211$  nA (n=3) in response to 1  $\mu\text{M}$  human NMU-25. Porcine NMU-25 (1  $\mu\text{M}$ ) also elicited a strong response ( $2150 \pm 330$  nA (n=3)) from oocytes injected with SNORF62 mRNA, but not in control oocytes (data not shown).



Detection of RNA coding for human SNORF62

RNA was isolated from multiple tissues (listed in Table 3) and assayed as described. Quantitative RT-PCR using a fluorogenic probe demonstrated RNA encoding human SNORF62 to be localized in highest abundance in peripheral organs, particularly in elements of the urogenital and gastrointestinal systems.

The highest levels are found in the testes. The uterus, prostate and kidney (both cortex and medulla) express SNORF62 RNA. This is consistent with functional studies and localization of NMU that are found in uterus and prostate (see Background). However, NMU has not been localized to the kidney and its function there is not known. However, the presence in the kidney may be associated with NMU's effects at increasing arterial blood pressure (in rats, Minamino et al., 1985b).

The gastrointestinal system also has considerable amounts of SNORF62 RNA. The stomach, small intestine (pooled) as well as the duodenum express SNORF62 RNA. This is consistent with the high levels of NMU in the GI tract found by radioimmunoassay (RIA) (Domin et al. 1987) in both myenteric and submucosal plexuses of the gut (Ballesta et al. 1988) and the postulated role of NMU as a potent constrictor of smooth muscle. SNORF62 RNA is also present in the pancreas at levels equivalent to that seen in other regions of the GI tract, but the role of the receptor in this tissue is not clear. It is not known if the SNORF62 receptors are found on the pancreatic islets, acinar cells or are present on vasculature within the gland. Sumi et al. (1987) demonstrated an increase in blood flow in the pancreas after administration of NMU suggesting a vascular localization or function.

Other tissues expressing SNORF62 RNA include the lung, trachea, adrenal gland, and mammary gland, with lower levels in skeletal muscle, and heart. This broad distribution implies a broad

regulatory or modulatory activity, perhaps at the level of smooth muscle contraction or secretagogue actions within these tissues. As discussed in the Background, NMU directly affects cells of the adrenal gland to alter secretion and may therefore act as a secretagogue in other tissues as well.

CNS structures express SNORF62 RNA but at levels much lower than those seen in peripheral organs. Within the CNS, SNORF62 RNA has been detected in highest abundance in the cerebellum, dorsal root ganglia, hippocampus and spinal cord. NMU-like immunoreactivity was identified in each of these regions (Domin et al. 1987). Within the CNS, it is found in levels that are 5 to 25-fold less than that found in peripheral organs. The role of SNORF62 RNA in the CNS is not clear, however its broad distribution is consistent with the broad distribution of NMU found in the brain (Domin et al. 1987). The presence of NMU as well as SNORF62 RNA in the spinal cord, dorsal root ganglia, and medulla oblongata implies a role in sensory transmission or modulation.

In summary, SNORF62 RNA is broadly distributed, with highest concentrations in gastrointestinal and urogenital systems. Levels within the CNS are fairly low. This distribution implies regulation/modulation of multiple systems. Some of the effects of the peripheral actions of SNORF62 may be mediated by the actions of NMU on smooth muscle, and its CNS distribution suggests a role in the modulation of sensory transmission.

#### Detection of RNA coding for human SNORF72

RNA was isolated from multiple tissues (listed in Table 3) and assayed as described. Quantitative RT-PCR using a fluorogenic probe demonstrated RNA encoding human SNORF72 to be localized in highest abundance in the CNS. The CNS regions expressing the highest levels of SNORF72 RNA include the medulla

oblongata, pontine reticular formation, spinal cord, and thalamus (Table 3). This distribution is highly suggestive of a role in sensory transmission or modulation and is in sharp contrast to the pattern seen with SNORF62, which has a distribution primarily in peripheral organs. The exception to this CNS/peripheral organ pattern are the testes, which express high levels of both SNORF62 and SNORF72 RNA.

The hippocampus, hypothalamus and cerebral cortex all express moderate-high levels of SNORF72 RNA. Other CNS structures expressing SNORF72 RNA include the amygdala and cerebellum. Dorsal root ganglia also express SNORF72 RNA albeit at substantially lower levels than those found in the spinal cord, but comparable to those found in the amygdala and cerebellum.

The expression pattern of SNORF72 RNA in the CNS is consistent with the hypothesis that its ligand, NMU, is a sensory transmitter/modulator. NMU is found in the spinal cord, dorsal root ganglia, and medulla oblongata using radioimmunoassay (Domin et al. 1987) and immunohistochemistry (Ballesta et al. 1988). Its presence in other regions including hippocampus, hypothalamus and cerebral cortex implies a modulatory role in multiple systems within the CNS.

Peripheral organs expressing SNORF72 RNA include the kidney (medulla), lung and trachea. The function of SNORF72 in the kidney may be different from SNORF62 despite the fact that NMU is possibly an endogenous ligand for both. SNORF62 RNA is found in equivalent amounts in both the cortex and medulla of the kidney. SNORF72 RNA is found primarily in the medulla, suggesting different physiological functions for these two receptors in the kidney. It is not known at this time which cells in the kidney express SNORF62 and/or SNORF72 RNA. The broad distribution in multiple peripheral organs (Table 3) implies a broad regulatory or modulatory activity, perhaps at

5

In summary, SNORF72 RNA is expressed in highest abundance in the CNS, particularly in structures associated with sensory transmission/or modulation. This localization suggests a role for SNORF72 in the central actions of NMU whereas the localization of SNORF62 RNA suggest a role for SNORF62 in the peripheral actions of this peptide.

**TABLE 3**

Summary of distribution of mRNA coding for human SNORF62 and human SNORF72.

5      *Data is expressed as % of the highest expressing tissue.*

Region	snorf62 % of maximum	snorf72 % of maximum	Potential applications
adipose tissue	<1	not detected	Obesity and metabolic disorders
adrenal gland	31.65	< 1	Regulation of metabolic steroids, disorders of the adrenal gland, regulation of epinephrine release
amygdala	5.90	9.58	Depression, phobias, anxiety, mood disorders
cerebellum	21.52	6.25	Motor coordination disorders
cerebral cortex	8.42	20.28	Cognition, sensory and motor integration disorders
dorsal root ganglia	14.68	9.33	Sensory transmission disorders, pain
duodenum	26.96	trace	Gastrointestinal disorders
heart	12.66	2.63	Cardiovascular disorders
hippocampus	13.92	43.61	Cognition/memory disorders
hypothalamus	10.13	40.58	Appetite/obesity, neuroendocrine regulation disorders
kidney, cortex	47.00	1.03	Hypertension, electrolyte balance disorders
kidney, medulla	33.33	16.42	Hypertension, electrolyte balance disorders
liver	not detected	not detected	Metabolic disorders
lung	48.86	17.08	Respiratory disorders, asthma
mammary gland	20.25	1.46	Lactation disorders

<b>medulla oblongata</b>	7.27	100	Sensory transmission/ integration disorders, pain, cardiovascular disorders, respiratory disorders,
<b>pancreas</b>	45.70	trace	Endocrine disorders, diabetes, pancreatitis
<b>pituitary</b>	3.44	< 1	Endocrine/neuroendocrine disorders
<b>pontine reticular formation</b>	3.61	92.50	Sleep disorders, sensory modulation and transmission disorders
<b>prostate gland</b>	27.09	1.79	Benign prostatic hyperplasia and male sexual dysfunction
<b>salivary gland</b>	2.84	not detected	Digestive disorders
<b>skeletal muscle</b>	13.67	< 1	Musculoskeletal disorders
<b>small intestine</b>	73.80	1.74	Gastrointestinal disorders
<b>spinal cord</b>	13.16	80.00	Analgesia, sensory modulation and transmission disorders, pain
<b>spleen</b>	1.25	not detected	Immune disorders
<b>stomach</b>	39.62	5.58	Gastrointestinal disorders
<b>testes</b>	100.00	85.00	Male reproductive disorders, regulation of steroid hormones
<b>thalamus</b>	7.48	47.83	Sensory integration disorders, pain
<b>trachea</b>	27.85	8.92	Respiratory disorders, asthma
<b>uterus</b>	34.43	4.20	Gestational disorders, dysmenorrhea, female sexual dysfunction

The identification of SNORF62 and SNORF72 as members of the family of NMU receptors is supported by a variety of experimental results. Both receptors are activated by full length NMU (rat, porcine and human) as well as NMU-8. In membranes prepared from SNORF62- or SNORF72-transfected cells,

[<sup>125</sup>I]rat NMU-23 and [<sup>125</sup>I]NMU-8 bind with high affinity. Human NMU-25 also demonstrates activation of SNORF62 expressed in *Xenopus* oocytes. Taken together these results indicate that SNORF62 and SNORF72 are functional NMU receptors. Differential localization of SNORF62 RNA predominantly to the periphery and SNORF72 RNA predominantly to the CNS suggest different roles for these receptors *in vivo*.

Chromosomal localization of human SNORF62 and SNORF72 receptor genes

The human SNORF62 gene maps to SHGC-33253 which is localized to chromosome 2q34-q37. SNORF 72 maps to SHGC-8848, which is localized to chromosome 5q31.1-q31.3.

**Detection of RNA coding for rat SNORF 72:** mRNA was isolated from multiple tissues and assayed as described (See Table 4).

Quantitative RT-PCR using a fluorogenic probe demonstrated mRNA coding for rat SNORF72 to be localized in highest abundance in the ovary and uterus. This is consistent with functional studies and localization of neuromedin U. Neuromedin U has been localized to the uterus (Domin, et al., 1987) and it has been shown to potently contract uterine smooth muscle (Minamino, et al., 1985a and 1985b)..

The stomach and the duodenum also express SNORF72 RNA. As described previously, this is consistent with the high levels of NMU in the GI tract found by radioimmunoassay (RIA) (Domin, et al., 1987). SNORF72 RNA is also expressed in substantial amounts in the urinary bladder. Taken together, this localization is consistent with the postulated role of NMU as

5 In the rat CNS highest levels of SNORF72 are expressed in the spinal cord and medulla oblongata. In the human, these also express high levels of SNORF72. The presence of SNORF72 RNA in the medulla and spinal cord is suggestive of a role in sensory transmission of modulation.

15

20



Table 4

Summary of distribution of mRNA coding for rat SNORF72 receptors

5 RNA encoding SNORF72r is expressed as % of highest expressing tissue: uterus

Tissue	qRT-PCR	Potential applications
	% of max	
adipose	1.26	metabolic disorders
10 adrenal cortex	trace	regulation of steroid hormones
adrenal medulla	trace	regulation of epinephrine release
cerebellum	0.81	motor coordination
cerebral cortex	3.39	Sensory and motor integration, cognition
dorsal root ganglia	2.72	sensory transmission
15 duodenum	4.33	gastrointestinal disorders
heart	0.14	cardiovascular indications
hippocampus	4.27	cognition/memory
hypothalamus	13.14	appetite/obesity, neuroendocrine regulation
kidney	0.07	electrolyte balance, hypertension
20 liver	0.08	diabetes
lung	4.84	respiratory disorders, asthma
medulla oblongata	5.07	analgesia, motor coordination
ovary	78.50	reproductive function
pancreas	1.44	diabetes, endocrine disorders
25 pituitary	trace	endocrine/neuroendocrine regulation

5

**REFERENCES**

Austin, C., et al., "Cloning and characterization of the cDNA encoding the human neuromedin U (NmU) precursor: NmU expression in the human gastrointestinal tract" *J Molecular Endocrinology* **14**: 157-169 (1995).

Ballesta J., et al., "Occurence and developmental pattern of neuromedin U-immunoreactive nerves in the gastrointestinal tract and brain of the rat" *Neuroscience* **25**: 797-816 (1988).

Benito-Orfila, M.A., et al., "The motor effects of neuromedin U on rat stomach *in vitro*" *Eur J Pharmacol* **193**: 329-333 (1991).

Bradford, M.M., "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding", *Anal. Biochem.* **72**: 248-254 (1976).

Brown D.R. and F.L. Quito, "Neuromedin U octapeptide alters ion transport in porcine jejunum" *Eur J Pharmacol* **155**:159-162 (1988).

Bush, et al., "Nerve growth factor potentiates bradykinin-induced calcium influx and release in PC12 cells" *J. Neurochem.* **57**: 562-574(1991).

Chu, Y.Y., et al., "Characterization of the rat A2a adenosine receptor gene", *DNA Cell Biol.* **15(4)**: 329-337 (1996).

Cimini V., et al., "Modulation of galanin nad neuromedin U-like immunoreactivity in rat corticotrophes after alteration of endocrine status" *Cell Tissue Res* **272**: 137-146 (1993).

5     Domin J., et al., "Neuromedin U - A study of its distribution  
in the rat" *Peptides* **8**:779-784 (1987).

0

5

0

5

0

Gundersen, C.B., et al., "Serotonin receptors induced by exogenous messenger RNA in *Xenopus* oocytes" *Proc. R. Soc. Lond. B. Biol. Sci.* **219(1214)**: 103-109 (1983).

5

0

5

0

5

0

Maggi, C.A., et al., "Motor response of the human isolated small intestine and urinary bladder to porcine neuromedin U-8" *Br J Pharmacol* **99**: 186-188 (1990).

Malendowicz, L.K., et al., "Effects of neuromedin U-8 on the rat pituitary-adrenocortical axis" *In Vivo* **7**:419-422 (1993).

Malendowicz, L.K., et al., "Effects of Neuromedin U (NMU)-8 on the rat hypothalamo-pituitary-adrenal axis. Evidence of a direct effect of NMU-8 on the adrenal gland" *Neuropeptides* **26**:47-53 (1994a).

Malendowicz, L.K., et al., "Effects of neuromedin U-8 on the secretory activity of the rat adtenal cortex: Evidence for an indirect action requiring the presence of the zona medullaris" *Res Exp Med* **194**:69-79 (1994b).

Milligan, G. and S. Rees, "Chimaeric G alpha proteins: their potential use in drug discovery" *Trends Pharmacol. Sci.* 20: 118-124 (1999).

Minamino, N., et al., "Neuromedins: Novel smooth muscle

stimulation peptides identified in porcine spinal cord"  
*Peptides* 6:245-248 (1985a).

Minamino, N., et al., "Neuromedin U-8 and U-25: Novel uterus  
stimulating and hypertensive peptide identified in porcine  
spinal cord" *Biochem Biophys Res Commun* 130: 1078-1085  
(1985b).

Nandha, K.A., et al. "Autoradiographic analysis of specific  
neuromedin U binding sites in the rat uterus and brain" *J. of*  
*Endocrinology* 140(Suppl.): Abstract No. P177 (1994).

Nandha K.A. and S.R. Bloom, "Neuromedin U - An overview"  
*Biomedical Research* 14(Suppl.3):71-76 (1993).

Nandha, K.A., et al., "Characterization of the rat uterine  
neuromedin U receptor." *Endocrinology* 133:482-486 (1993).

O'Harte, F., et al., "Isolation, structural characterization  
and pharmacological activity of dog neuromedin U" *Peptides*  
12:11-15 (1991).

Quick, M.W. and Lester, H.A., "Methods for expression of  
excitability proteins in *Xenopus* oocytes", *Meth. Neurosci.* 19:  
261-279 (1994).

Sakura, N., et al., "Structure-activity relationships of rat  
neuromedin U for smooth muscle contraction" *Chem Pharm Bull*  
(Tokyo) 39: 2016-2020 (1991).

Salon, J.A. and Owicki, J.A., "Real-time measurements of  
receptor activity: Application of microphysiometric techniques  
to receptor biology" *Methods in Neuroscience* 25: 201-224,  
Academic Press (1996).

Smith, K.E., et al., "Expression cloning of a rat hypothalamic galanin receptor coupled to phosphoinositide turnover." *J. Biol. Chem.* 272: 24612-24616 (1997).

5 Spurney, R.F., et al., "The C-terminus of the thromboxane receptor contributes to coupling and desensitization in a mouse mesangial cell line", *J. Pharmacol. Exp. Ther.* 283(1): 207-215 (1997).

10 Steel, J.H., et al., "Localization of 7B2, Neuromedin B, and neuromedin U in specific cell types of rat, mouse, and human pituitary, in rat hypothalamus, and in 30 human pituitary and extrapituitary tumors" *Endocrinology* 122:270-282 (1988).

15 Sumi, S., et al., "Effect of synthetic neuromedin U-8 and U-25, novel peptides identified in porcine spinal cord, on splanchnic circulation in dogs" *Life Sci* 41:1585-1590 (1987).

20 Takahashi, T., et al., "Rat brain serotonin receptors in *Xenopus* oocytes are coupled by intracellular calcium to endogenous channels." *Proc. Natl. Acad. Sci. USA* 84(14): 5063-5067 (1987).

25 Tian, W., et al., "Determinants of alpha-Adrenergic Receptor Activation of G protein: Evidence for a Precoupled Receptor/G protein State." *Molecular Pharmacology* 45: 524-553 (1994).

30 Underwood, D.J. et al., "Structural model of antagonist and agonist binding to the angiotensin II, AT1 subtype, G protein coupled receptor", *Chem. Biol.* 1(4): 211-221 (1994).